

PLANT-DERIVED NATURAL PRODUCTS FROM THE AMERICAN CONTINENT FOR THE CONTROL OF PHYTOPATHOGENIC FUNGI: A REVIEW

Marco Vinicio Ramirez-Mares^{1,*} and Beatriz Hernandez-Carlos²

¹Instituto de Recursos, Universidad del Mar, Puerto Ángel, Oaxaca 70902, México;

²Instituto de Agroindustrias, Universidad Tecnológica de la Mixteca, Huajuapán de León, Oaxaca 69000, México

*Corresponding author's e-mail: jvinicio2000@yahoo.com.mx; marcoram@angel.umar.mx

Various pathogenic microorganisms affect plants. However, most diseases are caused by fungi, which have an enormous reproductive capacity and possess metabolic mechanisms capable to develop resistance to commercial fungicides. Fungi produce mycotoxins on crops and this represents a considerable risk to human and animal health. The most important genera of phytopathogenic fungi of the American continent are: *Aspergillus*, *Alternaria*, *Botrytis*, *Colletotrichum*, *Fusarium*, *Phytophthora*, *Penicillium* and *Pythium*, among others. Different strategies have been used for phytopathogenic fungi control, mainly by using synthetic fungicides. Currently, a great deal of attention has been paid towards exploitation of higher plant products as an innovative alternative in crops protection. The use of plant derived natural products has many advantages, such as little or no harmful side effects, rare cases of resistance, long-term control and completely or substantially eliminates the use of synthetic fungicides. Several plants belonging to the Anacardiaceae, Asteraceae, Caprifoliaceae, Lamiaceae, Liliaceae, Papilionacea, Poaceae, Verbenaceae, families, have showed fungicide potential. Therefore, this review addresses occurrence of the most important phytopathogenic fungi and the current state of research regarding the potential of vegetable biodiversity in the development of plant derived natural products for the management of these pathogens in the American continent.

Keywords: Phytopathogenic fungi, biofungicides, mycotoxins, allelochemicals, crops, integrated pest management

INTRODUCTION

The total human population is steadily increasing. According to the latest estimates of the Population Reference Bureau (PRB, 2014), the global population reached 7,238 million in 2014 and is projected to increase to 9,683 million in 2050. With approximately 393 thousand births get added each day. The population of Northern America (Canada and United States) will increase from 353 million in 2014 to 444 million in 2050. In contrast, the population in Latin America and the Caribbean will increase from 618 million in 2014 to 773 million in 2050. Such demographic changes would have serious implications for the environment, economy, health and quality of life for the people living in the American Continent. With this population growth, demand will also increase on water, energy, arable land and biological resources to provide adequate food supplies. In 1995 one-third of cropland of the world (1.5 billion hectares) had been abandoned because erosion made it unproductive (Pimentel et al., 1995). Each year about 10 million hectares of fertile cropland are lost due to soil erosion (Pimentel, 2006). In addition, the demand of biofuel will rise from 115 billion L in 2013 to 135 billion L in 2019 (IEA, 2014), that will limit the amount of land available for food production (land-use change), thereby compromising food security, threatening biodiversity or limiting low-income farmers access to land. In order to provide food for the population which is in permanent augmentation is imperative to increase the

production and solve the political and logistical problems of food distribution (Zhang, 2009). This is the current challenge for agriculture. More recent the Food and Agricultural Organization (FAO) estimates, indicate that 47 million people living in Latin America and the Caribbean were unable to meet their dietary energy requirements in 2011-2013 (FAO, 2013). The decrease losses of food during pre-harvest and post-harvest, can provide an effective way to increase food availability and reduce the land need for its production (Kader, 2005). Food are very prone to microbial spoilage, infection by microorganisms (bacteria, viruses, fungi, nematodes, phytoplasmas and viroids) that cause postharvest decay can occur before the harvest and can remain latent until storage, when environmental condition favor their development. Most diseases and crop production losses are caused by phytopathogenic fungi which are able to infect any tissue at any stage of plant growth (Agrios, 2005). Plant pathogenic fungi are characterized by high complexity and diversity, these biological variability allow them to growth in very climatologically different environments, since dry zones until wet hot regions (Agrios, 2005). Moreover, fungal spores are more durable and long-lived than other microorganisms, can easily hitch rides on humans and as humans move around the world, fungi of all kinds are landing in new habitats. Recently, Dean et al., (2012) published the Top 10 fungal plant pathogen list, based on scientific/economic importance. The Top 10 list includes, in rank order, (1) *Magnaporthe oryzae*; (2) *Botrytis cinerea*; (3) *Puccinia* spp.; (4) *Fusarium graminearum*; (5)

Fusarium oxysporum; (6) *Blumeria graminis*; (7) *Mycosphaerella graminicola*; (8) *Colletotrichum* spp.; (9) *Ustilago maydis* and (10) *Melampsora lini*.

In fruits, vegetables and cereals, there is a wide variety of fungal genera causing quality problems related to organoleptic characteristics, appearance, nutritional value and limited shelf life (Yanes et al., 2012; Dean et al., 2012). In addition, the production of mycotoxins by several fungi is responsible for allergic or toxic disorders among humans and livestock (Díaz-Dellavalle et al., 2011). More than 300 fungal metabolites are reported to be toxic to animals and man (Galvano et al., 2001). Fungal proliferation and mycotoxins occur more frequently in regions with high temperatures, high moisture, flash floods and unseasonal rains during harvest. Beyond good horticultural and agronomic practices, growers often rely on pesticides and fertilizers. The use of pesticides is essential in agriculture production, the loss of cereals, vegetables and fruits from pest injury would reach 32%, 54% and 78% respectively, without pesticides application (Cai, 2008). Synthetic fungicides are still the most important components in the management of fungal plant diseases (Croplife, 2012). Even though the use of synthetic fungicides controls the seed borne fungi seeds in an effective and efficient way, enabling the plants to grow to their maximum genetic potential, they cannot be applied to grains due to pesticide toxicity (Harris et al., 2001). The usefulness of synthetic fungicides are increasingly limited owing to toxicity to non-target organisms, development of resistance by pathogens, residual toxicity problems, pathogen resurgence, environmental pollution and public concerns about the harmful effects on human health (Sharma and Meshram, 2006; Wu et al., 2010). Over the past few years there has been an increasing political pressure to remove the most hazardous fungicides from the market and to develop formulation with less negative impacts. Thus, it is believed (Gullino et al., 2000) that fungicides should be optimized under integrated pest management (IPM) and the best option is to use eco-friendly approaches.

Plant derived fungicides used as extracts, essential oils or pure allelochemicals have enormous potential against plant pathogenic fungi, because the secondary metabolism of plants has evolved to protect them against microbial pathogens (Benner et al., 1993; Mabrouk, 2012; Elamathi et al., 2012). Higher plants contain a wide spectrum of secondary metabolites such as tannins, coumarins, alkaloids, phenolic compounds, quinones and phytoalexins, which have antifungal activity (Kagale et al., 2004). Biofungicides possess the following advantages; strong specificity, degrade to nontoxic products within a few days (eco-friendly), they are not phytotoxic, can be modified to improve performance and enhance quality and rare cases of resistance (Siddiqui and Gulzar, 2003; Gupta and Dikshit, 2010). The source plants for the biofungicides are endemic of the developing countries in America (most of the Latin America and the Caribbean countries); in these countries the benefit will be great if the eco-friendly biofungicides are part of an IPM strategy.

Phytopathogenic Fungi Identified In The American Continent : Fungi is an extensive group of eukaryotic microorganisms (Garrido et al., 2012), it is considered that ninety percent of plant pathogens are fungi (Buckley, 2008). Most of the plants can be attacked by some kind of fungus (or more) and is also known that the same phytopathogenic fungus can infect one or more types of plants, even of different families (O'Donnell et al., 2000; Agrios, 2005). Plant pathogenic fungi show a complex life cycles, the reproduction is primarily via spores (one or a few cells), that could be produced by sexual and asexual reproduction stages (Pérez-Martin and de Sena-Tomas, 2011; Garrido et al., 2012; Phillipson, 2012). The phytopathogenic fungi spend most of their life cycle in the plant, which serves as host, and other part of their cycle in the soil, on plant debris found there (as saprophytes), although some only develop as parasites (Crous et al., 2009). Generally, fungal reproductive bodies are formed on the surface of tissues of the host plant (or very close), which causes that the spores are able to disperse quickly and easily (O'Donnell et al., 2000; Garrido et al., 2012). Diseased plants differ morphologically and physiologically from healthy individuals (Rostás et al., 2003). The effects produced by fungi on plants may be of local type; when they affect a small tissue, or general damage, which depends of the type of fungus and plant infected (Garrido et al., 2011; Phillipson, 2012). The damage caused by the fungus is primarily a death of infected tissue (Rostás et al., 2003); showed as 'green islands', which are spots of photosynthetically active tissue surrounded by chlorotic areas, as well as wilted leaves, give plants with an altered appearance (Hatcher et al., 1994; Phillipson, 2012). The pathogen can also cause atrophy of the complete plant or small part, and in other cases they may cause excessive growth or hypertrophy, such as; galls, warts or tumors on the roots (Hardham, 2001; Phillipson, 2012). For fungi that affect the root, or the vascular system of the plant, the production of a yellow color in the plant blight is backed. The above effects can cause atrophy of the plant, which eventually lead to the death of the host (D'Mello and Macdonald, 2003). A comprehensive knowledge on the nature of plant hosts and the manner in which the healthy plants get infected is essential, in order to check the infiltration of host plant environment by different fungal pathogens. The following properties appear to be hallmarks of phytopathogen fungi: 1) highly developed infection structures; 2) limited secretory activity, especially of lytic enzymes; 3) carbohydrate-rich and protein-containing interfacial layers that separate fungal and plant plasma membranes; 4) long-term suppression of host defense; and 5) haustoria, which are specialized hyphae for nutrient absorption and metabolism (Mendgen and Hahn, 2002). These pathogens also could cause disease (e.g., infections, allergies) in man, and produce toxins that affect plants, animals and humans (De Lucca, 2007). Mycotoxins are toxic substances produced mostly as secondary metabolites by filamentous fungi that grow on seeds, grains, and feed in the field, or in storage, five fungal genera produce these metabolites (*Penicillium*, *Aspergillus*, *Fusarium*, *Claviceps*

Table 1. Major mycotoxins prevalent in the American continent

Fungal species	Mycotoxin	Human health problem	References
<i>Alternaria alternata</i> , <i>A. arborescens</i> , <i>A. tenuissima</i> , <i>A. solani</i> , <i>Alternaria sp.</i> , <i>Alternaria spp.</i> , <i>A. tagetica</i> and <i>A. zinniae</i>	Belong to three structural classes: the tetramic acid derivative, tenuazonic acid (TA); the dibenzopyrone derivatives, alternariol (AOH), alternariol methylether (AME) and altenuene; and the perylene derivatives, the altertoxins. Solanapyrones and zinnolide. Zinniol derivatives.	AOH and AME are mutagenic and cytotoxic to mammalian cells, and are suspected to be carcinogenic. They are one of the most common airborne allergens, as well as being one causative agent of phaeohyphomycosis in immunocompromised patients. Possible toxic and foetotoxic effect. Genotoxic, mutagenic, carcinogenic and cytotoxic effects. Mutagenic and clastogenic in various in vitro systems. Might be responsible for esophageal cancer.	Greco et al., 2012; Gambale et al., 1976; Pollock et al., 1982; Biggs et al., 1993; Biggs, 1995; Gamboa-Angulo et al., 2000; Gamboa-Angulo et al., 2002; Andersen et al., 2008; Andrew et al., 2009; Lourenco et al., 2009; Alexander et al., 2011; Díaz Dellavalle et al., 2011; Somma et al., 2011; Edin, 2012; Pavón-Moreno et al., 2012; Benavidez Roza et al., 2013
<i>Aspergillus brasiliensis</i> , <i>A. costaricensis</i> , <i>A. flavus</i> , <i>A. lacticoffeatus</i> , <i>A. nidulans</i> and <i>A. niger</i> .	Chaetocin, penitrem A, and xanthocillin, tenuazonic acid, aflatoxin G2, aflatoxin B1 and B2, chaetoglobosin C, ochratoxin A and spinulosin.	Potential liver carcinogens in humans. Also, increases risk of hepatocellular carcinoma in patients with chronic hepatitis B virus. Induce allergic disorders, such as; allergic asthma, allergic rhinitis, allergic sinusitis, allergic bronchopulmonary aspergillosis (ABPA), and hypersensitivity pneumonitis. Patients with cystic fibrosis are often reported to have complication of ABPA.	Banerjee and Kurup, 1998; Wogan, 1999; Freire et al., 2000
<i>Bipolaris oryzae</i> and <i>B. sorokiniana</i>	Cochliquinones and ophiobolins.	Cochliquinones are antagonists of the Human Chemokine Receptor CCR5.	Manandhar et al., 1998; Yoganathan et al., 2004; Consolo et al., 2012
<i>Botryodiplodia theobromae</i>	Botryodiplodin, (+)-7-iso-jasmonic acid and several minor jasmonates.	Botryodiplodin induces DNA-protein cross-links in mammalian cells.	San Gupta et al., 1966; Moulé et al., 1982; Hertel et al., 1997; Ramezani et al., 2007
<i>Botrytis cinerea</i>	Botrydial.	No reported to date.	Latorre et al., 2002; Muñoz et al., 2002; Daoubi et al., 2006; Asadollahi et al., 2013
<i>Claviceps spp.</i>	Toxic alkaloids.	Toxic amounts of alkaloids leading to the disease known as ergotism in humans and other animals consuming them. The Holy Fire or St. Anthony's Fire is the primary vascular manifestation, causing burning pain and gangrene in the feet and hands.	Richard et al., 2003; Alm and Elvevag, 2013; Ayarragaray, 2014
<i>Fusarium avenaceum</i> , <i>F. culmorum</i> , <i>F. graminearum</i> , <i>F. decemcellulare</i> , <i>F. moniliforme</i> , <i>F. oxysporum</i> , <i>F. sambucinum</i> , <i>F. sporotrichum</i> , <i>F. spp.</i> , <i>F. solani</i> and <i>F. verticillioides</i>	Fumonisin and Zearalenone.	These toxins have been shown to be carcinogenic in rats and mice. Seem to be the most likely cause of human esophageal cancer. They are considered to be potential causes of immunosuppression in humans leading to secondary disease.	Richard et al., 2003; Prandini et al., 2009; Tittlemier et al., 2013b
<i>Penicillium expansum</i> and <i>P. herquei</i> .	Ochratoxin.	It is a genotoxic carcinogen by induction of oxidative DNA lesions coupled with direct DNA adducts via quinone formation.	Richard et al., 2003; Pfohl-Leskowicz and Manderville, 2007; Hope and Hope, 2012

Table 2. Important fungal phytopathogens in the American continent

Fungal species	Plant Disease	Country	References
<i>Alternaria alternata</i> , <i>A. arborescens</i> , <i>A. tenuissima</i> , <i>A. solani</i> , <i>Alternaria sp.</i> , <i>Alternaria spp.</i> , <i>A. tagetica</i> and <i>A. zinniae</i>	Fruit rot of blueberry is the most common and severe postharvest problem. Spoiled tomatoes. Early blight on potato. Fruit rot of apple, pear, melon, apricot, grapes, raisins, strawberry, olive, citrus fruits and dried figs. Citrus fruits, olives and several other fruits and vegetables.	Argentina, USA, Canada, Mexico and Brazil.	Greco et al., 2012; Gambale et al., 1976; Biggs et al., 1993; Biggs, 1995; Gamboa-Angulo et al., 2000; Gamboa-Angulo et al., 2002; Andrew et al., 2009; Alexander et al., 2011; Díaz Dellavalle et al., 2011; Somma et al., 2011; Pavón-Moreno et al., 2012; Benavidez Roza et al., 2013
<i>Aspergillus brasiliensis</i> , <i>A. costaricensis</i> , <i>A. flavus</i> , <i>A. lacticoffeatus</i> , <i>A. nidulans</i> and <i>A. niger</i> . <i>Bipolaris oryzae</i> and <i>B. sorokiniana</i>	Black and White pepper. Nut kernels. Black rot of grapes, which subsequently may result in ochratoxin A contamination of grapes, grapes by-products, dried vine fruits, coffee, and cocoa They cause major foliar disease and grain discoloration on cereals, such as; rice, maize and sorghum, to cause a brown leaf spot disease	Brazil, Costa Rica and Venezuela.	Freire et al., 2000; Perrone et al., 2011; Adjovi et al., 2014
<i>Blumeria graminis</i>	Also is called the grass powdery mildew fungus and causes serious damage to a variety of cereal crops.	USA, Canada and Mexico.	Wyand and Brown, 2003; Inuma et al., 2007; Sánchez-Martín et al., 2011
<i>Botryodiplodia theobromae</i>	Infection of cocoa beans.	USA.	San Gupta et al., 1966; Fessehatzion and Olutiola, 1987; Ramezani et al., 2007
<i>Botrytis cinerea</i>	Spoiled tomatoes. Also, is the most important disease of table grapes (<i>Vitis vinifera</i> L.)	Chile and Argentina.	Latorre et al., 2002; Muñoz et al., 2002; Asadollahi et al., 2013
<i>Cladosporium herbarum</i> <i>C. sp.</i>	Causes a Leaf Spot on Marshmarigold (<i>Caltha</i> spp.) are members of the buttercup family (Ranunculaceae) and inhabit stream banks and wet meadows in subalpine habitats of the Rocky and Cascade Mountains.	USA.	Lago et al., 2004; Rodríguez-Rajo et al., 2005; Johnson et al., 2008
<i>Claviceps</i> spp.	This organism is known as a replacement parasite in that it replaces plant structures with fungal tissue called ergots or sclerotia.	Argentina and USA.	Richard et al., 2003; Fisher et al., 2005
<i>Colletotrichum acutatum</i> , <i>C. coccodes</i> , <i>C. fragariae</i> , <i>C. graminicola</i> , <i>C. gloeosporioides</i> , <i>C. lindemuthianum</i> , <i>C. spp.</i> and <i>C. truncatum</i> . <i>Macrophomina phaseolina</i> .	Citrus postbloom fruit drop epidemics and legume species. Is the cause of anthracnose with black sunken lesion. Affect tomato, onions and dry bean cultures.	Brazil, Canada and USA.	Buchwaldt et al., 2004; Rodríguez-Rajo et al., 2005; Hernández-Silva et al., 2007; Ciampi-Guillardi et al., 2014; Yang et al., 2014
<i>Phytophthora infestans</i> , <i>P. plurivora</i> and <i>P. ramorum</i> . <i>Phymatotrichum omnivorum</i> .	The causal agent of the late blight of solanaceous plant species, which can devastate potato and tomato crops.	Mexico and USA.	Baird et al., 2010; Saleh et al., 2010; Schinke and Germani, 2012; Iqbal and Mukhtar, 2014
<i>Pythium sp</i> and <i>P. ultimum</i> .	Phymatotrichum (cotton or Texas) root rot is caused by the soil-borne fungus.	USA.	Del Castillo-Munera et al., 2013; Eyre et al., 2013; Martín et al., 2014; Schoebel et al., 2014
<i>Rhizoctonia solani</i> .	Causes a variety of diseases including seed rots and damping-off, root, stem and fruit rots, foliar blights and postharvest decay.	Canada and USA.	Gunasekaran and Ahuja, 1975; Uppalapati et al., 2010
<i>Fusarium avenaceum</i> , <i>F. culmorum</i> , <i>F. graminearum</i> , <i>F. decemcellulare</i> , <i>F. moniliforme</i> , <i>F. oxysporum</i> , <i>F. sambucinum</i> , <i>F. sporotrichum</i> , <i>F. spp.</i> , <i>F. solani</i> and <i>F. verticillioides</i>	Causes black scurf on potato tubers, stolon and root pruning, and stem canker on above ground plant parts. Black scurf is characterised by the presence of black sclerotia on the tuber surface which result in a significant tuber blemish.	Colombia.	Vaartaja, 1974; Pane et al., 2011; Taylor et al., 2002
<i>Penicillium expansum</i> and <i>P. herquei</i> .	They are causes of wilts and scab or blight diseases of small grains and ear rot. Affect grains such as wheat, barley, oats, and corn.	Argentina, Canada, Uruguay and USA.	Gonzalez-Vera et al., 2010; Ferrucho et al., 2013; Thangavel et al., 2014
	Is the primary cause of blue mold, a common postharvest fruit rot disease of apple.	Argentina and USA.	Richard et al., 2003; Sampietro et al., 2011; Poole et al., 2013; Tittlemier et al., 2013a; Umpierrez-Failache et al., 2013; Malbrán et al., 2014
			Li and Xiao, 2008; Finn et al., 2013; Rouissi et al., 2013

and *Alternaria*) (Anjorin et al., 2013). The most prevalent classes of mycotoxins are; aflatoxins, trichothecenes (e.g., deoxynivalenol, T-2 toxin), zearalenone, ochratoxins, and fumonisins (Table 1). When ingested, inhaled, or absorbed through skin, mycotoxins may reduce appetite and general performance, and cause sickness or death in humans (Tanaka et al., 2007; Kumar et al., 2008). The global occurrence of mycotoxins is considered to be a major risk factor, according to FAO (2012), 25% of the world's commodities are annually affected by known mycotoxins. Mycotoxins are formed in the field during the growing season; however, they also are formed or increased during harvest, drying, and storage (Richard et al., 2003). In the United States only, the mean economic annual costs of crop losses caused by mycotoxins are estimated to be 932 million USD (Richard et al., 2003). During the last decade, the problem of these pathogenic fungal infections in American Continent, has reached epidemic proportions (O'Donnell et al., 2000; Bruin and Sabelis, 2010; Barres et al., 2012). The classification of plant pathogenic fungi considers the host range and disease symptoms produced, but a problem arises when morphologically similar fungi produce similar disease symptoms, but have no clearly defined host range (Taylor and Pedras, 1995). In this review, we present a compilation of the most important phytopathogenic fungi in our continent (Table 2).

Vegetable Extracts With Antifungal Activity : A total of 143 plants from America, belonging to 43 botanical families (109 genera) possess toxic activity against phytopathogenic fungi species of 28 genera (Table 3). These pathogens attack crops, stored grains, seeds and food products, and include species of *Alternaria*, *Aspergillus*, *Bipolaris*, *Botryodiplodia*, *Botrytis*, *Cladosporium*, *Colletotrichum*, *Corynespora*, *Curvularia*, *Drechslera*, *Epicoccum*, *Exserobolus*, *Fusarium*, *Gaeumannomyces*, *Geotrichum*, *Gerlachia*, *Leptosphaeria*, *Macrophomina*, *Monilia*, *Moniliophthora*, *Mycosphaerella*, *Neurospora*, *Penicillium*, *Pestalotia*, *Phaeoisariopsis*, *Phomopsis*, *Pyricularia*, *Pythium*, *Rhizopus*, *Sarocladium*, *Sclerotinia*, *Sclerotium*, *Sitophilus*, *Sporothrix*, *Trichoderma* and *Trichophyton* genera. The botanical families with the highest number of species with antifungal activity are Asteraceae (22), Fabaceae (18), Lamiaceae (13), Rutaceae (9), Verbenaceae (8), Euphorbiaceae (8) and Poaceae (7). There are 61 plants fungicides against *Colletotrichum* genera, followed by 42 plants with toxic activity against *Fusarium* and *Alternaria* species, while only one toxic vegetal specie for the following genera is described; *Drechslera*, *Epicoccum*, *Gaeumannomyces*, *Leptosphaeria*, *Monilia*, *Mycosphaerella*, *Neurospora*, *Pestalotia*, *Sitophilus*, *Trichoderma*, and *Trichophyton*. Plant treatments for fungicide tests were mixtures of essential oils (Jaramillo et al., 2012; Landero-Valenzuela et al., 2013), or methanol, ethanol, dichloromethane, chloroform, and aqueous extracts. Some authors considered alternatives to the practical application of treatments, using non-common solvent extracts as cocoa butter and lanolin (Castillo et al., 2010) or

added to chitosan edible films (Avila-Sosa et al., 2012). In addition, treatment of plant or extracts to improve the antifungal activity was used, such as the fermentation of aqueous extracts of *Flourensia cernua* (De León et al., 2013), which were most effective against *F. oxysporum* than those prepared in ethanol (Jasso de Rodríguez et al., 2007). In addition, the abiotic elicitation (UV light) of *Brassica oleracea* was used to produce the toxic caulilexins A-C to *Leptosphaeria maculans*, *Rhizoctonia solani*, and *Sclerotinia sclerotiorum* (Pedras et al., 2006).

Compounds With Fungicide Activity Essential oils: Gas chromatography-mass spectrometry analysis of mixtures of essential oils with fungicide activity show as main compounds: 1,8-cineole, 4-carene, 5-methylfurfural, benzene acetaldehyde, borneol, isocaryophyllene, carvacrol, caryophyllene, citral, elemol, germacrane D, germacrene, hexadecanoic acid, limonene, *m*-cymene, methylcinnamate, methyl-eugenol, *n*-nonanal, *p*-cymene, sabinene, safrole, spathulenol, thymol, *trans*-cinnamaldehyde, α -bisabolol, α -cadinol, α -terpinene, α -terpinyl acetate, β -myrcene, β -phellandrene, 5-phenyl-1,3-pentadiene, capillarin, and β -pinene (Table 3). However, some authors showed the fungicide activity of isolated compounds, such as 5-methylfurfural, benzene acetaldehyde, *n*-nonanal, and carvacrol, all actives against *Colletotrichum* species (Kobaisy et al., 2001; Demirci et al., 2006). Also, it was described the synergistic fungicidal activity of compounds present in the essential oil mixture from *Chrysanthamnus nauseosus* against *Botrytis cinerea*, with more than 50 identified compounds, where non-major component showed this toxic activity (Tabanca et al., 2007).

Other Compounds with Fungicide Activity: Although Asteraceae is the family most studied, from Fabaceae species have been isolated more fungicides compounds, different to that found in essential oils (Figure 1). Diterpenoid 1 (Cotoras et al., 2004) and bicyclic sesquiterpenic alcohol (2) are toxic to *Botrytis* (Ortiz-Núñez et al., 2007). Other terpenes are 15,16-dihydroxypimar-8(14)-en-3-one (3), fungicide against *Colletotrichum gloeosporioides* (Peraza-Sánchez et al., 2005) and the sesquiterpenes polygodial (4), drimenol, drimenin, and isodrimenol that inhibited the growth of *Gaeumannomyces graminis* (Monsálvez et al., 2010). Compound 1 produces permeabilization of the cell membrane of *B. cinerea* through the induction of *p*-nitrophenyl butyrate esterase and laccase production (Cotoras et al., 2004). Twelve bioactive alkaloids (Figure 1) were described; among various structures, the amide type 6 and 12, resulted toxic to *Sclerotium* and *Botrytis* species respectively (Meepagala et al., 2010; Ramírez-Chávez et al., 2000). Quinoline alkaloids as 7, 11, 2-*n*-nonyl-4-quinolone, and 13 were active against *Alternaria*, *Epicoccum*, *Pestalotia*, *Drechslera*, *Fusarium*, *Colletotrichum* and *Botrytis* species (Oliva et al., 2003; Li et al., 2005). While those of the pyrrol type as caulexin A (5), B and C were toxic to *Leptosphaeria*, *Rhizoctonia*, *Sclerotinia* (Pedras et al., 2006); and quinolizidine alkaloids

Table 3. Vegetable species from America with antifungal activity

Family	Specie	Patogen	Observations	References	Country
Amaranthaceae	<i>Chenopodium ambrosioides</i>	<i>Fusarium oxysporum</i>	Growth inhibition by essential oils (α -terpinene, <i>p</i> -cymene, 4-carene)	Jaramillo et al., 2012	Colombia
	<i>Teloxys ambrosioides</i> (<i>Chenopodium ambrosioides</i>)	<i>Fusarium sp.</i>	Growth inhibition by essential oils	Barrera-Necha and García-Barrera, 2008	Mexico
Amaryllidaceae	<i>Allium sativum</i>	<i>Drechslera tritici-repentis</i> , <i>Bipolaris sorokiniana</i>	Garlic extract and allicin protect wheat seed infected	Perelló et al., 2013	Argentina
		<i>Colletotrichum gloeosporioides</i>	Growth, germination and sporulation inhibition of essential oils <i>in vitro</i> and <i>in vivo</i> assays	Landero-Valenzuela et al., 2013	Mexico
Anacardiaceae	<i>Myracrodruon urundeuva</i>	<i>Fusarium decemcellulare</i> , <i>F. monoliforme</i> , <i>F. oxysporum</i> , <i>F. solani</i>	Growth inhibition by extracts (gallic acid, flavonoids, luteolin, cinnamic derivatives, proanthocyanidins, hydrolysable tannins, and leucoanthocyanidins)	Sá et al., 2009	Brazil
Asparagaceae	<i>Agave lechuguilla</i>	<i>Colletotrichum gloeosporioides</i> , <i>Penicillium digitatum</i> , <i>Rhizoctonia solani</i>	500-5000 μ L/L produces 90% sporulation inhibition 500-3000 μ L/L >55% growth inhibition	Jasso de Rodríguez et al., 2011	Mexico
	<i>Yucca carnerosana</i>	<i>Rhizopus stolonifer</i> , <i>Colletotrichum gloeosporioides</i> , <i>Penicillium digitatum</i>	Extracts with polyphenols showed growth inhibition (CI ₅₀ 32.5 ppm) Growth inhibition 3000-5000 μ L/L 80% 3000 μ L/L sporulation inhibition	Castillo-Reyes et al., 2015 Jasso de Rodríguez et al., 2011	Mexico
	<i>Yucca filifera</i>	<i>Rhizopus stolonifer</i> , <i>Colletotrichum gloeosporioides</i> , <i>Penicillium digitatum</i>	Growth inhibition 2000 μ L/L Sporulation inhibition 1000 μ L/L	Jasso de Rodríguez et al., 2011	Mexico
Apocynaceae	<i>Allamanda blanchetti</i>	<i>Fusarium sp.</i> , <i>Cladosporium sp.</i> , <i>Curvularia sp.</i> , <i>Alternaria sp.</i>	Ethanol extracts protects <i>Pterogyne nitens</i> seeds at 100 and 500 ppm	Ferreira-Medeiros, et al., 2013	Brazil
	<i>Calotropis procera</i>	<i>Fusarium solani</i> , <i>Colletotrichum gloeosporioides</i> , <i>Neurospora sp.</i>	Growth inhibition 20,100 Da-Protein IC ₅₀ 32.1-67 μ g/mL	Teixeira de Freitas et al., 2011	Brazil
Asparagaceae	<i>Furcraea cahum</i>	<i>Alternaria tagetica</i> , <i>Colletotrichum gloeosporioides</i>	Growth inhibition by ethanol extract	Gamboa-Angulo et al., 2008	Mexico
	<i>Bonellia flammea</i>	<i>Curvularia verruculosa</i> , <i>C. lunata</i> , <i>Exserobolium rostratum</i> , <i>Corynespora cassiicola</i> ,	Aqueous extracts showed growth mycelial growth >50%, sporulation inhibition >58% (<i>C. verruculosa</i> , <i>E. rostratum</i> , <i>C. lunata</i>) and germination inhibition >73% (<i>C. verruculosa</i> , <i>E. rostratum</i> , <i>C. lunata</i> , <i>C. cassiicola</i>)	Moo-Koh et al., 2014	Mexico
Asteraceae	<i>Adenophyllum aurantium</i>	<i>Fusarium solani</i> , <i>Alternaria alternata</i>	Sporulation and mycelial growth inhibition (MIC 7.78-27 mg/mL) by methanol and ethyl acetate extracts from roots and aerial parts.	Lira de León et al., 2014	Mexico
	<i>Ageratum conyzoides</i>	<i>Aspergillus flavus</i>	Essential oils at > 0.1 μ g/mL inhibited mycelial growth and aflatoxins production	Nogueira et al., 2010	Brazil
	<i>Ambrosia hispida</i>	<i>Alternaria tagetica</i> , <i>Colletotrichum gloeosporioides</i>	Growth inhibition by ethanol extract	Gamboa-Angulo et al., 2008	Mexico
	<i>Alloispermum integrifolium</i>	<i>Fusarium solani</i> , <i>Alternaria alternata</i>	Sporulation inhibition (MIC 16.88-33.76 mg/mL) by methanol extracts from aerial parts	Lira de León et al., 2014	Mexico
	<i>Arnica longifolia</i>	<i>Colletotrichum acutatum</i> , <i>C. fragariae</i> , <i>C.</i>	Growth inhibitor by essential oils (carvacrol and α -bisabolol)	Tabanca et al., 2007	USA

Phytochemicals and plant extracts from America against phytopathogenic fungi

<i>Artemisia annua</i>	<i>gloeosporioides</i> , <i>Alternaria alternata</i>	Growth inhibition <i>in vitro</i> but not <i>in vivo</i> by methanol extract	Carvalho et al., 2011	Brazil
<i>Artemisia dracunculus</i>	<i>Colletotrichum fragariae</i> , <i>C. gloeosporioides</i> , <i>C. accutatum</i>	Growth inhibition by essential oils (Methyl-eugenol)	Meepagala et al., 2002	USA
<i>Artemisia ludoviciana</i>	<i>Colletotrichum lindemuthianum</i>	Growth and spore germination inhibition (borneol, spathulenol, derivatives of caryophyllene and spathulenol)	Damián-Badillo et al., 2008	Mexico
<i>Aster hesperius</i>	<i>Colletotrichum accutatum</i> , <i>C. fragariae</i> , <i>C. gloeosporioides</i> , <i>Alternaria tagetica</i>	Growth inhibition by essential oils (hexadecanoic acid, carvacrol)	Tabanca et al., 2007	USA
<i>Calea urticifolia</i>		Growth inhibition by ethanol extract	Gamboa-Angulo et al., 2008	Mexico
<i>Conyza canadensis</i>	<i>Colletotrichum acutatum</i> , <i>C. fragariae</i> , <i>C. gloeosporioides</i>	Growth inhibition by (4Z,8Z)-matricaria lactone (43)	Queiroz et al., 2012	USA
<i>Chrysactinia mexicana</i>	<i>Aspergillus flavus</i>	Growth inhibition by essential oils	Cárdenas-Ortega et al., 2005	Mexico
<i>Chrysothamnus nauseosus</i>	<i>Colletotrichum acutatum</i> , <i>C. fragariae</i> , <i>C. gloeosporioides</i>	Growth inhibition by essential oils (β -phellandrene and β -pinene)	Tabanca et al., 2007	USA
<i>Cynara scolymus</i>	<i>Alternaria spp</i>	Growth inhibition by aqueous extract at MIC 5 μ g/mL and saline buffer and acid extracts at MIC 10 μ g/mL	Díaz-Dellavalle et al., 2011	Uruguay
<i>Fluorensia cernua</i>	<i>Fusarium oxysporum</i> , <i>Penicillium expansum</i> <i>Rhizoctonia solani</i>	Fermented extract produces growth inhibition at 500 ppm Cocoa butter extract produces growth inhibition CI ₅₀ 420 ppm Extracts with polyphenols showed growth inhibition (CI ₅₀ 16.3 ppm)	De León et al., 2013 Castillo et al., 2010 Castillo-Reyes et al., 2015	Mexico Mexico Mexico
	<i>Fusarium oxysporum</i> , <i>Alternaria sp.</i> , <i>Rhizoctonia solani</i>	Total growth inhibition at $\leq 1500 \mu$ l/l by ethanol extract	Jasso de Rodríguez et al., 2007	Mexico
<i>F. retinophylla</i>	<i>Fusarium oxysporum</i> , <i>Alternaria sp.</i> , <i>Rhizoctonia solani</i>	Total growth inhibition at $\leq 1500 \mu$ l/l by ethanol extract	Jasso de Rodríguez et al., 2007	Mexico
<i>Haplopappus Greenei</i>	<i>Colletotrichum acutatum</i> , <i>C. fragariae</i> , <i>C. gloeosporioides</i>	Growth inhibition by essential oils (carvacrol)	Demirci et al., 2006	USA
<i>Heliopsis longipes</i>	<i>Sclerotium cepivorum</i> , <i>S. rolfsii</i>	Growth inhibition by alkamide affinin (17)	Ramírez-Chávez et al., 2000	Mexico
	<i>Colletotrichum lindemuthianum</i> , <i>Sporothrix schenckii</i>	Growth and germination inhibition by chloroform and methanol extracts	Damián-Badillo et al., 2008	Mexico
<i>Mutisia friesiana</i>	<i>Cladosporium cucumerinum</i>	Moderate growth inhibition by 5-methylcoumaranones (30-33)	Vituro et al., 2004	Argentina
<i>Pseudognaphalium vira vira</i>	<i>Botrytis cinerea</i>	Growth inhibition by 3 β -hydroxy-kaurenoic acid (3)	Cotoras et al., 2004	Chile
<i>Tagetes erecta</i>	<i>Rhizoctonia solani</i> Kühn	Growth inhibition by aqueous extract at 0.5 g/mL	Espejo-Quispe et al., 2010	Cuba
<i>T. lucida</i>	<i>Colletotrichum lindemuthianum</i> , <i>Sporothrix schenckii</i>	Growth and germination inhibition by chloroform and methanol extracts	Damián-Badillo et al., 2008	Mexico
Balantiopsaceae	<i>Balantiopsis cancellata</i>	Growth inhibition by Phenylethyl-2- <i>trans</i> - β -methylthioacrylate (44)	Labbe et al., 2005	Chile
Brassicaceae	<i>Brassica oleracea</i> var. botrytis	Growth inhibition by caulilexins A-C (11-13)	Pedras et al., 2006	Canada
	<i>Leptosphaeria maculans</i> , <i>Rhizoctonia solani</i> , <i>Sclerotinia sclerotium</i> <i>Alternaria alternata</i>	Spore germination inhibition with broccoli juice at 0.07, 0.11 and 0.15 μ g/ μ L	Flores-Córdova et al., 2013	Mexico
	<i>Colletotrichum gloeosporioides</i>	Glucoraphanin inhibited sporulation (IC ₅₀ 0.65 μ g/ μ L)	Lara-Viveros et al., 2014	Mexico

	<i>Raphanus raphanistrum</i> L.	<i>C. acutatum</i> , <i>F. oxysporum</i> , <i>B.cinerea</i>	Ethanol extracts showed moderate fungistatic effect on growth and germination	Sánchez-León, et al., 2015	Colombi a
Boraginaceae	<i>Tournefortia densiflora</i>	<i>Fusarium solani</i> , <i>Alternaria alternata</i>	Sporulation and mycelial growth inhibition (MIC 11.75-16.3 mg/mL) by methanol extracts from aerial parts and roots	Lira de León et al., 2014	Mexico
Cactaceae	<i>Opuntia ficus-indica</i>	<i>Colletotrichum gloeosporioides</i> , <i>Fusarium descencelulare</i> , <i>F. lateritium</i> , <i>F. moniliforme</i> , <i>F. oxysporum</i> and <i>F. solani</i> . <i>Rhizoctonia solani</i>	Growth inhibition by extract and lectin Extracts with polyphenols showed growth inhibition (CI ₅₀ 39.4 ppm)	Santana et al., 2009 Castillo-Reyes et al., 2015	Brazil Mexico
Caricaceae	<i>Carica papaya</i>	<i>Fusarium spp.</i>	Growth inhibition by ethanol extract at 0.625 mg/mL	Chávez-Quintal, et al., 2011	Mexico
Cleomaceae	<i>Cleome viscosa</i>	<i>Alternaria solani</i>	Growth and germination inhibition by ethanol extracts at 2 mg/mL	Pupo-Blanco et al., 2011	Cuba
Clusiaceae	<i>Garcinia mangostana</i> L.	<i>Botrytis cinerea</i>	Ethanol extracts from peels of fruits showed protection of cauliflower	Angel et al., 2014	Colombi a
Combretaceae	<i>Terminalia catappa</i>	<i>Rhizoctonia solani</i> Kühn	Growth inhibition by aqueous extract at <0.65 mg/mL	Espejo-Quiste et al., 2008	Cuba
Convolvulaceae	<i>Ipomoea murucoides</i>	<i>Fusarium solani</i> , <i>Alternaria alternata</i>	Sporulation inhibition (MIC 5.77-11.54 mg/mL) by ether extract from leaves	Lira de León et al., 2014	Mexico
Cornaceae	<i>Camptotheca acuminata</i>	<i>Alternaria alternata</i> , <i>Epicoccum nigrum</i> , <i>Pestalotia guepinii</i> , <i>Dreschlera sp.</i> , <i>Fusarium avenaceum</i>	Growth inhibition by camptothecin (14), trifolin (35) and hyperoside (36) at < 50µg/mL	Li et al., 2005	USA
Cucurbitaceae	<i>Momordica charantia</i>	<i>Fusarium sp.</i> , <i>Cladosporium sp.</i> , <i>Curvularia sp.</i> , <i>Alternaria sp.</i>	Ethanol extracts protects <i>Pterogyne nitens</i> seeds at 100, 500 and 1000 ppm	Ferreira-Medeiros, et al., 2013	Brazil
Cupressaceae	<i>Juniperus lucayana</i>	<i>Botrytis cinerea</i>	Growth inhibition by widdrol (4) at 50 ppm	Ortiz-Nuñez et al., 2007	Cuba
Euphorbiaceae	<i>Acalypha cuspidata</i>	<i>Fusarium solani</i> , <i>Alternaria alternata</i>	Sporulation inhibition (MIC 12.5 mg/mL) by methanol extracts from aerial parts	Lira de León et al., 2014	Mexico
	<i>A. diversifolia</i>	<i>Fusarium solani</i>	Growth inhibition by dichloro-methane extract at 1 mg/mL	Niño et al., 2012	Colombi a
	<i>A. gaumeri</i>	<i>Alternaria tagetica</i> , <i>Colletotrichum gloeosporioides</i> , <i>Fusarium oxysporum</i> , <i>Rhizopus sp.</i>	Growth inhibition by ethanol extract	Gamboa-Angulo et al., 2008	Mexico
	<i>A. subviscida</i>	<i>Alternaria alternata</i>	Sporulation inhibition (MIC 8.51 mg/mL) by methanol extracts from leaves	Lira de León et al., 2014	Mexico
	<i>Croton chichinensis</i>	<i>Alternaria tagetica</i> , <i>Colletotrichum gloeosporioides</i> , <i>Fusarium oxysporum</i> , <i>Rhizopus sp.</i>	Growth inhibition by ethanol extract (Alkaloids)	Gamboa-Angulo et al., 2008	Mexico
	<i>Euphorbia antisiphilitica</i>	<i>Colletotrichum gloeosporioides</i> , <i>Rhizoctonia solani</i> , <i>Fusarium oxysporum</i> , <i>Alternaria alternata</i>	Growth inhibition by ellagitannin- (34) rich ethanolic fraction (MW 860.7 umas)	Ascacio-Valdés et al., 2013	Mexico
	<i>Euphorbia sp</i>	<i>Fusarium solani</i>	Growth inhibition by dichloro-methane extract at 1 mg/mL	Niño et al., 2012	Colombi a

Phytochemicals and plant extracts from America against phytopathogenic fungi

Fabaceae	<i>Acacia penulata</i>	<i>Colletotrichum gloeosporioides</i>	Growth, sporulation and germination inhibition by 15,16-dihydroxypimar-8(14)-en-3-one (5)	Peraza-Sánchez et al., 2005	México
	<i>Anadenanthera colubrina</i>	<i>Alternaria alternata</i>	Growth inhibition <i>in vitro</i> but not <i>in vivo</i>	Carvalho et al., 2011	Brazil
	<i>Caesalpinia cacalaco</i> Bonpl.	<i>Colletotrichum lindemuthianum</i>	Growth and germination inhibition by phenolic compounds rich extract	Veloz-García et al., 2010	Mexico
	<i>Calia secundiflora</i>	<i>Alternaria solani</i> , <i>Fusarium oxysporum</i> , <i>Monila fruticola</i> , <i>Curvularia lunata</i> , <i>Penicillium herquei</i> , <i>Trichoderma pseudokoningii</i> ,	Growth inhibition by extracts containing quinolizidine alkaloids (cytosine, 15)	Pérez-Láinez et al., 2008	Mexico
	<i>Canavalia ensiformis</i>	<i>Curvularia lunata</i> , <i>Penicillium herquei</i> , <i>Trichoderma pseudokoningii</i> ,	Growth and germination inhibition at sub-micromolar concentration by an urease	Becker-Ritt et al., 2007	Brazil
	<i>Crotalaria pallida</i>	<i>Fusarium oxysporum</i>	Growth inhibition by a 5340 Da peptide	Pelegrini et al., 2009	Brazil
	<i>Dioclea guianensis</i> Benth	<i>Colletotrichum gloeosporioides</i>	Germination inhibition by a ConA-like lectin	Araujo-Filho et al., 2010	Brazil
	<i>Geoffroea decorticans</i> Burkart	<i>Botrytis cinerea</i>	Germination and growth of conidia	Boiteux et al., 2015	Argentina
	<i>Gliricidia sepium</i>	<i>Colletotrichum acutatum</i>	Coumarin (0.25 and 0.5%) showed fungicide activity while ortho coumaric acid at 05 % showed fungistatic effect	Urdaneta et al., 2015	Venezuela
	<i>Glycine max</i>	<i>Curvularia lunata</i> , <i>Penicillium herquei</i> , <i>Trichoderma pseudokoningii</i> ,	Growth and germination inhibition at sub-micromolar concentrations by an urease (IC ₅₀ < 0.81µM)	Becker-Ritt et al., 2007	Brazil
	<i>Luetzelburgia auriculata</i>	<i>Colletotrichum lindemuthianum</i> , <i>Fusarium solani</i> , <i>Aspergillus niger</i>	Growth inhibition by a 123.5 kDa-lectin	Melo et al., 2005	Brazil
	<i>Lupinus mexicanus</i>	<i>Rhizoctonia solani</i> , <i>Sclerotium solfsii</i> , <i>Fusarium oxysporum</i>	Growth inhibition by alkaloid extract (lupanine, 16)	Zamora-Natera et al., 2008	Mexico
	<i>Phaseolus vulgaris</i> L.	<i>C. lindemuthianum</i> , <i>Phaeoisariopsis griseola</i>	Methanol-water extracts from leaves showed growth inhibition	Noreña-Ramírez et al., 2014	Colombia
	<i>Piscidia piscipula</i>	<i>Colletotrichum gloeosporioides</i>	Growth inhibition by methanol extract	Peraza-Sánchez et al., 2005	Mexico
	<i>Pithecellobium dulce</i>	<i>Colletotrichum gloeosporioides</i>	Growth inhibition (by methanol extract)	Peraza-Sánchez et al., 2005	Mexico
	<i>Pongamia glabra</i>	<i>Sclerotinia sclerotiorum</i>	Growth inhibition by oils in combination with neem oils	Garcia et al., 2012	Brazil
Juglandaceae	<i>Vigna unguiculata</i> (L.) Walp	<i>C. lindemuthianum</i> , <i>Phaeoisariopsis griseola</i>	Methanol-water extracts from leaves showed growth inhibition	Noreña-Ramírez et al., 2014	Colombia
	<i>Zuccagnia punctata</i>	<i>Phomopsis longicolla</i> , <i>Colletotrichum truncatum</i>	Growth inhibition by chloroform, and ethanol extracts (MIC 62.5-100 µg/mL) Chalcones (39 , 40), flavanone (42) and caffeoyl ester (41)	Svetaz et al., 2004	Argentina
	<i>Carya illinoensis</i>	<i>Phytium sp</i> , <i>Colletotrichum truncatum</i> , <i>C. coccodes</i> , <i>Alternaria alternata</i> , <i>Fusarium solani</i> , <i>F. vertillioides</i> , <i>F. sambucinum</i> , <i>Rhizoctonia solani</i>	Growth inhibition by polyphenolic extracts at 0.20 mg/L	Osorio et al., 2010	Mexico
Lamiaceae	<i>Hedeoma multiflora</i>	<i>Aspergillus section Flavi</i>	Growth and germination inhibition and reduction of aflatoxins by essential oils	Bluma et al., 2008	Argentina
	<i>Origanum vulgare</i>	<i>Colletotrichum lindemuthianum</i>	Growth and germination inhibition by methanol extract	Andrade-Pinto et al., 2010	Brazil

		<i>Aspergillus niger</i> , <i>A. flavus</i>	Germination and growth inhibition by essential oils in combination with those of <i>Rosmarinus officinalis</i> L.	Lima de Sousa et al., 2013	Brazil
	<i>Ocimum gratissimum</i>	<i>Colletotrichum lindemuthianum</i>	Growth and germination inhibition by methanol extract	Andrade-Pinto et al., 2010	Brazil
	<i>Rosmarinus officinalis</i> L.	<i>Alternaria spp</i>	Growth inhibition by acid extract at MIC 1.25µg/mL	Díaz-Dellavalle et al., 2011	Uruguay
	<i>R. officinalis</i>	<i>Alternaria alternata</i>	Growth inhibition of aqueous extract in combination with <i>L. alba</i> ones.	Tagami et al., 2009	Brazil
	<i>Salvia macrochlamys</i>	<i>Colletotrichum acutatum</i> , <i>C. fragariae</i> , <i>C. gloeosporioides</i> , <i>Alternaria spp</i>	Growth inhibition of essential oils (Carvacrol, Thymol)	Tabanca et al., 2007	USA
	<i>S. officinalis</i>	<i>Alternaria spp</i>	Growth inhibition by acid extract at MIC 1.25µg/mL	Díaz-Dellavalle et al., 2011	Chile
	<i>S. recognita</i>	<i>Colletotrichum acutatum</i> , <i>C. fragariae</i> , <i>C. gloeosporioides</i> , <i>Alternaria spp</i>	Growth inhibition of essential oils (Carvacrol, Thymol)	Tabanca et al., 2007	USA
	<i>S. sclarea</i>	<i>Alternaria spp</i>	Growth inhibition by acid extract at MIC 1.25µg/mL and saline buffer extract at MIC 5µg/mL	Díaz-Dellavalle et al., 2011	Chile
	<i>Satureja macrostema</i>	<i>Colletotrichum lindemuthianum</i> , <i>Sporothrix schenckii</i>	Growth and germination inhibition by chloroform and methanol extracts	Damián-Badillo et al., 2008	Mexico
	<i>Thymus vulgaris</i>	<i>Fusarium oxysporum</i>	Growth inhibition by thyme essential oil	Barrera-Necha et al., 2008	Mexico
	<i>T. vulgaris</i>	<i>Colletotrichum gloeosporioides</i> , <i>Rhizopus stolonifer</i>	Growth inhibition of essential oils <i>in vitro</i> and <i>in vivo</i> assays	Bosquez-Molina et al., 2010	Mexico
		<i>Colletotrichum graminicola</i>	Growth inhibition of aqueous extract	Tagami et al., 2009	Brazil
	<i>Vitex gaumeri</i>	<i>Alternaria tagetica</i>	Growth inhibition by ethanol extract	Gamboa-Angulo et al., 2008	Mexico
Lauraceae	<i>Cinnamomum zeylanicum</i>	<i>Colletotrichum gloeosporioides</i>	Growth, germination and sporulation inhibition of essential oils <i>in vitro</i> and <i>in vivo</i> assays	Landero-Valenzuela et al., 2013	Mexico
		<i>Fusarium oxysporum</i> , <i>Alternaria alternata</i> , <i>Geotrichum candidum</i> , <i>Trichoderma spp.</i> , <i>Penicillium digitatum</i> , <i>Aspergillus niger</i>	Growth mycelial inhibition by aqueous extracts (MIC 500 ppm). Cinnamaldehyde and kaempferol were identified as Main constituent	Cáceres Rueda de León et al., 2013	Mexico
	<i>C. verum</i>	<i>Penicillium digitatum</i>	Growth inhibition by 0.5 % essential oils applied into chitosan films	Ávila-Sosa et al., 2012	Mexico
	<i>Ocotea quixos</i> (Lam.)	<i>Phytium ultimum</i>	Growth inhibition by essential oils (trans-cinnamaldehyde, methylcinnamate, 1,8-cineole)	Bruni et al., 2004	Ecuador
Malvaceae	<i>Guazuma ulmifolia</i>	<i>Sclerotium cepivorum</i>	Hexane and dichloromethane extracts showed growth mycelial inhibition >84% at 10000 ppm	Ramírez-Aguayo, et al., 2015	Mexico
	<i>Hibiscus cannabinus</i>	<i>Colletotrichum fragariae</i> , <i>C. gloeosporioides</i> , <i>C. accutatum</i>	Growth inhibition by essential oil (5-methylfurfural, benzene acetaldehyde and <i>n</i> -nonanal)	Kobaisy et al., 2001	USA
	<i>Malva sp.</i>	<i>Gerlachia oryzae</i>	Growth inhibition of aqueous extract	Knaak et al., 2013	Brazil
Meliaceae	<i>Trichila minutiflora</i>	<i>Rhizopus sp.</i>	Growth inhibition by ethanol extract	Gamboa-Angulo et al., 2008	Mexico
Melastomataceae	<i>Miconia argyrophyla</i>	<i>Colletotrichum lindemuthianum</i>	Growth and germination inhibition by methanol extract	Andrade-Pinto et al., 2010	Brazil
Musaceae	<i>Musa acuminata</i>	<i>Mycosphaerella fijiensis</i>	Growth inhibition by phenalenone compounds (45, 46)	Otálvaro et al., 2007	Colombia
Myrtaceae	<i>Corymbia citriodora</i>	<i>Pyricularia (Magnaporthe) grisea</i> <i>Aspergillus spp.</i>	Germination and sporulation total inhibition at 0.15 mg/mL of essential oil and citronellal, which	Aguiar et al., 2014	Brazil

	<i>Eugenia caryophyllata</i>	<i>Colletotrichum musae</i> <i>Fusarium oxysporum</i> , <i>Alternaria alternata</i> , <i>Geotrichum candidum</i> , <i>Trichoderma</i> spp., <i>Penicillium digitatum</i> , <i>Aspegillus niger</i>	is main component (61.78 %) Growth mycelial inhibition by aqueous extracts (MIC 750-800 ppm). Gallic acid and kaempferol are main constituents.	Cáceres Rueda et al., 2013	Mexico
	<i>Melaleuca quinquenervia</i> (Cav)	<i>Sarocladium oryzae</i> , <i>Bipolaris oryzae</i>	Growth inhibition (100%) essential oil	Duarte et al., 2014	Cuba
	<i>Melaleuca alternifolia</i>	<i>Macrophomina phaseolina</i> , <i>Sclerotinia sclerotiorum</i> , <i>Alternaria alternata</i>	Growth inhibition by essential oils at 0.2 %	Martins et al., 2010	Brazil
	<i>Melaleuca quinquenervia</i> (Cav)	<i>Sarocladium oryzae</i> , <i>Bipolaris oryzae</i>	Growth inhibition (100%) essential oil	Duarte et al., 2014	Cuba
	<i>Myrcia fallax</i>	<i>Colletotrichum lindemuthianum</i>	Growth and germination inhibition by methanol extract	Andrade-Pinto et al., 2010	Brazil
	<i>Syzygium aromaticum</i>	<i>Fusarium oxysporum</i>	Growth inhibition by essential oil	Barrera-Necha et al., 2008	
		<i>Sclerotium cepivorum</i>	Growth inhibition by aqueous extract at 1%	Montes-Belmont et al., 2006	Mexico
		<i>Aspergillus</i> section Flavi	Growth and germination inhibition and reduction of aflatoxins by essential oils	Bluma et al., 2008	Argentina
Piperaceae	<i>Piper aduncum</i> L.	<i>Sclerotinia sclerotiorum</i>	Growth inhibition by aqueous extract	Garcia et al., 2012	Brazil
	<i>Piper aduncum</i> subsp. <i>ossanum</i>	<i>Curvularia lunata</i> , <i>Sarocladium oryzae</i> <i>Bipolaris oryzae</i>	Growth inhibition (100%) essential oil	Duarte et al., 2014	Cuba
	<i>P. auritum</i>	<i>Curvularia lunata</i> , <i>Sarocladium oryzae</i> <i>Bipolaris oryzae</i>	Growth inhibition (100%) essential oil	Duarte et al., 2014	Cuba
		<i>Colletotrichum acutatum</i> , <i>C. gloeosporioides</i> , <i>Botrydiploida theobromae</i>	Growth inhibition by <i>n</i> -hexane extract and essential oil (Safrole, apiol)	Pineda et al., 2012	Colombia
	<i>P. holtonii</i>	<i>Colletotrichum acutatum</i> , <i>C. gloeosporioides</i> , <i>Botrydiploida theobromae</i>			
Poaceae	<i>Cymbopogon nardus</i>	<i>Pyricularia (Magnaporthe) grisea</i> <i>Aspergillus</i> spp. <i>Colletotrichum musae</i>	Germination and sporulation inhibition at 0.15 mg/mL of essential oil and citronellal, which is main component (36.6 %)	Aguiar et al., 2014	Brazil
	<i>Cymbopogon citratus</i>	<i>Gerlachia oryzae</i>	Growth inhibition by aqueous extract	Knaak et al., 2013	Brazil
		<i>Alternaria alternata</i> , <i>Bipolaris</i> sp.	Growth inhibition by extract (IC ₅₀ 75.83 µg/mL) and citral (IC ₅₀ 58.24 µg/mL)	De Lima Guimarães et al., 2011	Brazil
	<i>C. nardus</i>	<i>Macrophomina phaseolina</i>	Growth inhibition by essential oils at 1.5%	Sánchez-García et al., 2008	Cuba
	<i>Hordeum vulgare</i> L.	<i>Fusarium culmorum</i>	<i>Fusarium</i> proteinase inhibition by proteins	Pekkarinen et al., 2003	USA
	<i>Oryza sativa</i>	<i>Aspergillus flavus</i>	Growth and aflatoxins production inhibition by phenolic extracts	Dos Santos Oliveira et al., 2008	Brazil
	<i>Triticum aestivum</i>	<i>Aspergillus flavus</i>	Growth and aflatoxins production inhibition by phenolic extracts	Dos Santos Oliveira et al., 2008	Brazil
	<i>Zea mays</i>	<i>Alternaria</i> spp	Growth inhibition by aqueous extract at MIC 5µg/mL	Díaz-Dellavalle et al., 2011	Chile
Phytolaccaceae	<i>Phytolacca</i>	<i>Colletotrichum</i>	Conidial production and growth	Hernández et	Argentina

	<i>tetramera</i>	<i>gloeosporioides</i>	inhibition by ethanol and aqueous extracts	al., 2013	a
	<i>Petiveria alliaceae</i>	<i>Gerlachia oryzae</i>	Growth inhibition by aqueous extract	Knaak et al., 2013	Brazil
Primulaceae	<i>Jacquinia macrocarpa</i>	<i>Fusarium verticillioides</i>	A fraction from a butanol partition showed growth inhibition (100%)	Valenzuela-Cota et al., 2014	Mexico
Rubiaceae	<i>Galium mexicanum</i>	<i>Fusarium solani</i> , <i>Alternaria alternata</i>	Sporulation inhibition (MIC 15 mg/mL) by methanol extracts from aerial parts	Lira de León et al., 2014	Mexico
Rutaceae	<i>Amyris texana</i>	<i>Botrytis cinerea</i>	Growth inhibition by a chromene amide (10)	Meepagala et al., 2010	USA
	<i>Citrus aurantifolia</i>	<i>Colletotrichum gloeosporioides</i> , <i>Rhizopus stolonifer</i> <i>Aspergillus nidulans</i> , <i>Colletotrichum acutatum</i>	Growth inhibition of essential oils <i>in vitro</i> and <i>in vivo</i> assays Conidia survival reduction by coumarins (22, 23, 26, 27) and furocoumarins (24, 25, 28, 29) under photo treatment	Bosquez-Molina et al., 2010 De Menezes et al., 2014	Mexico Brazil
	<i>C. sinensis</i>	<i>Bipolaris oryzae</i>	Growth inhibition (100%) essential oil	Duarte et al., 2014	Cuba
	<i>Ruta graveolens</i> L.	<i>Colletotrichum acutatum</i> , <i>Botrytis cinerea</i>	Growth inhibition by furanocoumarins and alkaloids (19 , 20 , 21)	Oliva et al., 2003	USA
	<i>Zanthoxylum fagara</i>	<i>Colletotrichum acutatum</i> , <i>Fusarium oxysporum</i> ,	Growth inhibition by essential oils (germacrene, elemol and α -cadinol)	Prieto et al., 2011	Colombia
	<i>Z. rhoifolium</i>	<i>Fusarium oxysporum</i> , <i>Colletotrichum acutatum</i>	Growth inhibition by essential oils (β -myrcene, β -phellandrene and germacrene D)	Prieto et al., 2011	Colombia
	<i>Z. monophyllum</i>	<i>Fusarium oxysporum</i> <i>Sitophilus oryzae</i> , <i>Colletotrichum acutatum</i>	Growth inhibition by essential oils (sabinene, 1.8-cineole and α -cadinol)	Prieto et al., 2011	Colombia
Sapindaceae	<i>Matayba elaeagnoides</i>	<i>Colletotrichum lindemuthianum</i>	Growth and germination inhibition by methanol extract	Andrade-Pinto et al., 2010	Brazil
Siparunaceae	<i>Siparuna arianeae</i>	<i>Colletotrichum lindemuthianum</i>	Growth and germination inhibition by methanol extract	Andrade-Pinto et al., 2010	Brazil
Solanaceae	<i>Capsicum annum</i> <i>var. glabriusculum</i>	<i>Alternaria alternata</i> , <i>Fusarium oxysporum</i>	Conidial germination inhibition (85.3-96%) of phenolic and carotene extracts from fruits	Rodriguez-Maturino et al., 2015	Mexico
	<i>Solanum lycopersicum</i>	<i>Moniliophthora perniciosa</i>	Growth inhibited by alfa-tomatine isolated from leaves	Gomes et al., 2014	Brazil
	<i>Solanum tuberosum</i> L-	<i>Aspergillus flavus</i>	Mycelial growth and aflatoxins production inhibition by phenolic compounds	Dos Santos Oliveira et al., 2008	Brazil
Styracaceae	<i>Styrax pohlilii</i>	<i>Colletotrichum lindemuthianum</i>	Growth and germination inhibition by methanol extract	Andrade-Pinto et al., 2010	Brazil
Verbenaceae	<i>Lantana achyranthifolia</i>	<i>Fusarium moniliforme</i>	Growth inhibition by essential oils (carvacrol, isocaryophyllene) at IC ₅₀ 100-180µg/mL	Hernández et al., 2008	Mexico
		<i>F. solani</i> , <i>Alternaria alternata</i>	Sporulation inhibition (MIC 6.25-12.5 mg/mL) by methanol extract from leaves	Lira de León et al., 2014	Mexico
	<i>L. camara</i>	<i>Alternaria solani</i>	Growth and germination inhibition of ethanol extracts	Pupo-Blanco et al., 2011	Cuba
		<i>Bipolaris oryzae</i>	Growth inhibition by aqueous extract	Knaak et al., 2013	Brazil
	<i>Lippia alba</i>	<i>Alternaria spp</i>	Growth inhibition by acid and saline extract at MIC 25 µg/mL	Díaz-Dellavalle et al., 2011	Chile
	<i>L. dulcis</i>	<i>Alternaria solani</i>	Growth and germination inhibition of ethanol extracts	Pupo-Blanco et al., 2011	Cuba
	<i>L. graveolens</i>	<i>Aspergillus niger</i> , <i>Trichophyton</i>	Growth inhibition by essential oils (carvacrol, α -terpinyl acetate, <i>m</i> -	Hernández et al., 2008	Mexico

		<i>mentagrophytes</i> , <i>Rhizoctonia solani</i> , <i>Fusarium sporotrichum</i> , <i>F. moniliforme</i> , <i>Aspergillus niger</i> <i>Rhizopus stolonifer</i> , <i>Colletotrichum</i> <i>gloeosporioides</i> , <i>Penicillium digitatum</i> <i>Aspergillus niger</i>	cymene) at IC ₅₀ 10-90µg/mL		
	<i>L. berlandieri</i> <i>Schauer</i> <i>L. berlandieri</i>	<i>Fusarium oxysporum</i> , <i>Alternaria alternata</i> , <i>Geotrichum candidum</i> , <i>Trichoderma spp.</i> , <i>Penicillium digitatum</i> , <i>Aspergillus niger</i>	Growth and sporulation inhibition of ethanol extract at 500-3000µl/L	Jasso de Rodríguez et al., 2011	Mexico
	<i>L. turbinata</i> var <i>Integrifolia</i>	<i>Aspergillus section Flavi</i>	Growth inhibition by 0.5 % essential oils applied into chitosan films	Ávila-Sosa et al., 2012	Mexico
			Growth mycelial inhibition by aqueous extracts (MIC 200-250 ppm). Gallic acid, carvacrol, thymol and caryophyllene were identified as main constituents	Cáceres Rueda de León et al., 2013	Mexico
			Growth and germination inhibition and reduction of aflatoxins by essential oils	Bluma et al., 2008	Argentina
Winteraceae	<i>Drimys winteri</i> J.R. Forst. & G. Forst.	<i>Gaeumannomyces graminis</i>	Growth inhibition by essential oil (α-pinene,β-pinene, limonene, β-myrcene) and <i>n</i> -hexane extract containing polygodial (6), drimenol (7), drimenin (8), isodrimenol (9)	Monsálvez et al., 2010	Chile
Xanthorrhoeaceae	<i>Aloe vera</i>	<i>Alternaria spp.</i>	Growth inhibition by acid extract at MIC 25 µg/mL	Díaz-Dellavalle et al., 2011	Chile
Zingiberaceae	<i>Zingiber officinale</i>	<i>Bipolaris oryzae</i> , <i>Gerlachia oryzae</i>	Growth inhibition of aqueous extract and essential oil	Knaak et al., 2013	Brazil
Zygophyllaceae	<i>Larrea cuneifolia</i>	<i>Fusarium oxysporum</i> , <i>Penicillium notatum</i> , <i>Aspergillus niger</i>	Growth inhibition by ethanol extract	Quiroga et al., 2001	Argentina
	<i>Larrea divaricata</i>	<i>Fusarium graminearum</i> , <i>F. solani</i> , <i>F. verticillioides</i> , <i>Macrophomina phaseolina</i> <i>Penicillium notatum</i> , <i>Aspergillus niger</i> , <i>Fusarium oxysporum</i>	Growth inhibition by chloroform extract and phenolic compounds (38) MIC 12.5-250 µg/mL	Vogt et al., 2013	Mexico
	<i>L. tridentata</i>	<i>Phytium sp.</i> , <i>Colletotrichum truncatum</i> , <i>C. coccodes</i> , <i>Alternaria alternata</i> , <i>Fusarium solani</i> , <i>F. verticillioides</i> , <i>F. sambucinum</i> , <i>Rhizoctonia solani</i> <i>R. solani</i> <i>R. solani</i>	Growth inhibition by ethanol extract Growth inhibition by mono and polyphenolics rich extracts at 0.20 mg/L Extracts with polyphenols showed growth inhibition (CI ₅₀ 58.4 ppm) Lanolin extract produces growth inhibition (CI ₅₀ 185 ppm)	Quiroga et al., 2001 Osorio et al., 2010 Castillo-Reyes et al., 2015 Castillo et al., 2010	Argentina Mexico Mexico Mexico

(8, 9) were toxic to *Monilia*, *Alternaria*, *Fusarium* and *Rhizoctonia* species (Pérez-Láinez et al., 2008; Zamora-Natera et al., 2008). Coumarins as 4-hydroxycoumarin, 7-hydroxycoumarin, 7-methoxycoumarin (14), 8-methoxypsoralen, citropten (15), 5-methoxypsoralen (16), and isopimpinellin (Figure 1) were described with antifungal activity against *Colletotrichum* and *Aspergillus* (De Menezes et al., 2014), and the closely related multisicoumaranones A (17), B, C and D, resulted toxic to *Cladosporium* species (Viturro et al., 2004). Phenolic and

polyphenolic compounds (Figure 1) of twelve vegetable species have shown broad-spectrum fungicide activity against eleven fungus genera, the active compounds isolated are ellagitannin (18), trifolin (19), hyperoside (20), lignans (21, 22), chalcones (23, 24), a caffeic acid derivative (25) and flavanone (26). About antifungal mechanisms, it was reported that the compound 22 is a fungal endo-β-(1,3)-glucanase inhibitor (Vargas-Arispuru et al., 2009). Others active compounds (Figure 1) are acetylenic lactone (27) (Queiroz et al., 2012), 2-phenylethanol ester (28) (Labbe et




Figure 1. Structures of compounds of vegetable origin with fungicide activity identified in the American continent

al., 2005), and the unusual 2-hydroxy-1-*H*-phenalen-one (29) and 2-methoxy-1-*H*-phenalen-one, which are fungicides against *Colletotrichum*, *Cladosporium*, and *Mycosphaerella*, respectively (Otálvaro et al., 2007). Other compound also toxic to *Colletotrichum* was the known glucosinolate glucoraphanin, isolated from *Brassica oleracea* (Lara-Viveros et al., 2014)

Plants are able to produce proteinaceous compounds, between those described with fungicide activity are lectins (glycoproteins that agglutinate and immobilize micro-organisms) and microbial proteinase inhibitors (inhibit

enzymes that degrade plant cell walls). Examples of lectins and fungal proteinase inhibitors with toxic activity on phytopathogen fungus are those isolated from *Dioclea guianensis* (Araújo-Filho et al., 2010), *Luetzelburgia auriculata* (Melo et al., 2005), *Calotropis procera* (De Freitas et al., 2011) and *Hordeum vulgare* (Pekkarinen and Jones, 2003) (Table 3). In addition, the binding of ConA-like lectin with ungerminated conidia of *Colletotrichum gloeosporioides* inhibit the germination (Araújo-Filho et al., 2010). But when the lectin interact with outer portion of cell wall and inner surface, the synthesis of fungal cell or cell

wall structure and function are disrupted. Such is the case of growth inhibition by agglutinin (LAA) isolated from *Lutzelburgia auriculata*, which inhibited the fungal growth and provoked interference on the transport of intracellular protons to the external medium (Melo et al., 2005). Other proteinaceous compounds are ureases; nickel-dependent metalloenzymes that catalyze the hydrolysis of urea to ammonia and carbon dioxide, as those described from *Glycine max*, which produced plasmolysis and cell wall injuries during the growth of *Curvularia lunata*, *Penicillium herquei* or *Trichoderma pseudokoningii* (Becker-Ritt et al., 2007).

CONCLUSIONS

The development of organic agriculture as projected by FAO (2009) may increase the use of biofungicides where few alternative pesticides are accessible. Even though that the eco-friendly fungicides are more pertinent for use in industrialized countries for the organic food production, they can play a much important role in the protection of food products in developing countries. Nowadays, there are more than 100 kinds of biopesticides in the world (Xu, 2008). The countries that use more biopesticides in the world are Mexico, Canada and the United States (44% global production), while Latin America and the Caribbean accounts for 9% (Qin and Kong, 2006). The developing countries are the source of endemic plants and this may have a great impact in improving food safety, through the production of biopesticides to promote food security. The majority of the evidences of fungitoxicity of plant extracts presented in this review were supported from *in vitro* evaluations, it is imperative to test the bioactive extracts on farm investigations under natural infections against a wide range of pathogens. The challenges to commercial application of plant extracts as biofungicides include the availability of plant material, efficacy at least equivalent to the current pesticide used, not toxicity against crop plants and that no compromise the quality of the crops. Moreover, as the phytochemicals profile of plant species can vary depending on climatic, geographic or genetic factor, among others, biofungicides manufacturers must to ensure that their products have a consistent and standardized composition. As the secondary metabolites occur in mixtures in the crude extracts, a good option is the guide-isolation and purification of the bioactive fungicide compounds, in order to elucidate their mechanisms of action, improve their effectiveness and enhance their production in the plant. In summary, sustainable food security cannot rely on the use of synthetic pesticides, endemic resources of biopesticides must be utilized in the production and post harvest protection of food products in developing countries, in order to ensure that the populations will have access to sufficient and healthy food.

REFERENCES

Adjovi, Y.C.S., S. Bailly, B.J.G. Gnonlonfin, S. Tadrast, A. Querin, A. Sanni, I.P. Oswald, O. Puel and J.D. Bailly.

2014. Analysis of the contrast between natural occurrence of toxigenic *Aspergilli* of the Flavi section and aflatoxin B1 in cassava. Food Microbiol. 38:151-159.
- Agrios, G.N. 2005. Plant Pathology. 5th ed. San Diego, CA: Elsevier Academic Press.
- Aguiar, R.W.D.S., M.A. Ootani, S.D. Ascencio, T.P. Ferreira, M.M.D. Santos and G.R.D. Santos. 2014. Fumigant antifungal activity of *Corymbia citriodora* and *Cymbopogon nardus* essential oils and Citronellal against three fungal species. Sci. World J. I.D. 492138, 8.
- Alexander, J., D. Benford, A. Boobis, S. Ceccatelli, B. Cottrell, J. Cravedi, A. Di Domenico, D. Doerge, E. Dogliotti and L. Edler. 2011. Scientific opinion on the risks for animal and public health related to the presence of *Alternaria* toxins in feed and food. EFSA J. 9:2407-2504.
- Alm, T., and B. Elvevag. 2013. Ergotism in Norway. Part 2: The symptoms and their interpretation from the eighteenth century onwards. Hist. Psychiatr. 24:131-147.
- Andersen, B., A. Dongo and B.M. Pryor. 2008. Secondary metabolite profiling of *Alternaria dauci*, *A. porri*, *A. solani*, and *A. tomatophila*. Mycol. Res. 112:241-250.
- Andrade Pinto, J.M., E.A. Souza and D.F. Oliveira. 2010. Use of plant extracts in the control of common bean anthracnose. Crop Prot., 29:838-842.
- Andrew, M., T.L. Peever and B.M. Pryor. 2009. An expanded multilocus phylogeny does not resolve morphological species within the small-spored *Alternaria* species complex. Mycologia. 101:95-109.
- Ángel, A.D., J.D.D. Flórez and M.P.C. González. 2014. Evaluación de la actividad fungistática del extracto de la cáscara del mangostino (*Garcinia mangostana* L.) en *Botrytis cinerea* para la biopreservación de la coliflor (*Brassica oleracea* var. botrytis). Alimentos Hoy. 22: 59-80.
- Anjorin, T.S., E.A. Salako and H.A. Makun. 2013. Control of Toxigenic Fungi and Mycotoxins with Phytochemicals: Potentials and Challenges, p 181-202. In: . D.H. Makun (ed.) Mycotoxin and Food Safety in Developing Countries. InTech
- Araújo-Filho, J.H., I.M. Vasconcelos, A.S. Martins-Miranda, D.M.F. Gondim and J.T.A. Oliveira. 2010. A ConA-like Lectin from *Dioclea guianensis* Benth. has antifungal activity against *Colletotrichum gloeosporioides*, unlike its homologues, ConM and ConA. J. Agric. Food Chem., 58:4090-4096.
- Asadollahi, M., E. Fekete, L. Karaffa, M. Flipphi, M. Árnýasi, M. Esmaeili, K.Z. Váczy and E. Sándor. 2013. Comparison of *Botrytis cinerea* populations isolated from two open-field cultivated host plants. Microbiol. Res. 168:379-388.
- Ascacio-Valdés, J., E. Burboa, A.F. Aguilera-Carbo, M. Aparicio, R. Pérez-Schmidt, R. Rodríguez and C.N. Aguilar. 2013. Antifungal ellagitannin isolated from *Euphorbia antisiphilitica* Zucc. Asian Pac. J. Trop. Biomed. 3:41-46.

- Avila-Sosa, R., E. Palou, M.T. Jiménez-Munigua, G.V. Nevarez-Moorillon, A.R. Navarro-Cruz and A. López-Malo. 2012. Antifungal activity by vapor contact of essential oils added to amaranth, chitosan or starch edible films. *Int. J. Food Microbiol.* 153:66-72.
- Ayarragaray, J.E. 2014. Ergotism: a change of perspective. *Ann. Vasc. Surg.* 28:265-268.
- Baird, R. E., P.A. Wadl, T. Allen, D. McNeill, X. Wang, J.K. Moulton, T.A. Rinehart, H.K. Abbas, T. Shier and R.N. Trigiano. 2010. Variability of United States isolates of *Macrophomina phaseolina* based on simple sequence repeats and cross genus transferability to related genera within botryosphaeriaceae. *Mycopathologia.* 170:169-180.
- Banerjee, B., and V.P. Kurup. 1998. Molecular biology of *Aspergillus allergens*. *Immunol. Allergy Clin. North Am.* 18:601-618.
- Barrera-Necha, L.L., and L.J. García Barrera. 2008. Actividad antifúngica de aceites esenciales sobre el crecimiento de *Fusarium* sp. aislado de papaya (*Carica papaya*). *Revista UDO Agrícola.* 8:33-41.
- Barrera-Necha, L.L., C. Garduño-Pizaña and L.J. García Barrera. 2009. In vitro Antifungal activity of essential oils and their compounds on mycelial growth of *Fusarium oxysporum* f. sp. Gladioli (Massey) Snyder and Hansen. *Plant Pathol. J.* 8:17-21.
- Barres, B., J. Carlier, M. Seguin, C. Fenouillet, C. Cilas and V. Ravigne. 2012. Understanding the recent colonization history of a plant pathogenic fungus using population genetic tools and Approximate Bayesian Computation. *Heredity.* 109:269-279.
- Becker-Ritt, A.B., A.H.S. Martinelli, S. Mitidieri, V. Feder, G.E. Wassermann, L. Santi, M.H. Vainstein, J.T.A. Oliveira, L.M. Fiuza, G. Pasquali and C.R. Carlini. 2007. Antifungal activity of plant and bacterial ureases. *Toxicon.* 50: 971-983.
- Benavidez Roza, M.E., A. Patriarca, G. Cabrera and V.E. Fernández Pinto. 2013. Determinación de perfiles de producción de metabolitos secundarios característicos de especies del género *Alternaria* aisladas de tomate. *Rev. Iberoam. Micol.* 31:119-124.
- Benner, J.P. 1993. Pesticidal Compounds from Higher Plants. *Pest. Sci.* 39:95-102.
- Biggs, A.R., M. Ingle and W.D. Solihati. 1993. Control of *Alternaria* infection of fruit of apple cultivar Nittany with calcium chloride and fungicides. *Plant Dis.* 77:976-980.
- Biggs, A.R. 1995. Detection of latent infections in apple fruit with paraquat. *Plant Dis.* 79:1062-1067.
- Bluma R., M.R. Amaiden and M. Etcheverry. 2008. Screening of Argentine plant extracts impact on growth parameters and aflatoxin B1 accumulation by *Aspergillus* section Flavi. *Int. J. Food Microbiol.* 122:114-125.
- Boiteux, J.J., M.V. Hapon, M.D.L.Á. Fernández, G.S. Lucero and P.H. Pizzuolo. 2015. Efecto del extracto acuoso de chañar (*Geoffroea decorticans* Burkart) sobre *Botrytis cinerea*, como posible alternativa para su control durante poscosecha de uva de mesa. *Revista de la Facultad de Ciencias Agrarias. Universidad Nacional de Cuyo,* 47:241-250.
- Bosquez-Molina, E., E. Ronquillo-de Jesús, S. Bautista-Baños, J.R. Verde-Calvo and J. Morales-López. 2010. Inhibitory effect of essential oils against *Colletotrichum gloeosporioides* and *Rhizopus stolonifer* in stored papaya fruit and their possible application in coatings. *Postharvest. Biol. Tec.* 57:132-137.
- Bruin, J. and M. Sabelis. 2010. Trends in Acarology: Proceedings of the 12th International Congress. Springer.
- Bruni, R., A. Medici, E. Andreotti, C. Fantin, M. Muzzoli, M. Dehesa, C. Romagnoli and G. Sacchetti. 2004. Chemical composition and biological activities of Ishpingo essential oil, a traditional Ecuadorian spice from *Ocotea quixos* (Lam.) Kosterm. (Lauraceae) flower calices. *Food Chem.* 85: 415-421.
- Buchwaldt, L., K.L. Anderson, R.A. Morrall, B.D. Gossen and C.C. Bernier. 2004. Identification of Lentil Germ Plasm Resistant to *Colletotrichum truncatum* and Characterization of Two Pathogen Races. *Phytopathology.* 94:236-243.
- Buckley, M. 2008. The fungal kingdom: Diverse and essential roles in earth's ecosystem. In: AAM Colloquium. The Fungal Kingdom: Diverse and Essential Roles in Earth's Ecosystem. Tucson, Arizona, USA: American Society for Microbiology.
- Cai, D.W. 2008. Understand the role of chemical pesticides and prevent misuses of pesticides. *Bull. Agric. Sci. Technol.* 1:36-38.
- Cáceres Rueda de León, I., R. Colorado Vargas, E. Salas Muñoz, L.N. Muñoz Castellanos and L. Hernández Ochoa. 2013. Actividad Antifúngica in vitro de Extractos Acuáticos de Especies contra *Fusarium oxysporum*, *Alternaria alternata*, *Geotrichum candidum*, *Trichoderma* spp., *Penicillium digitatum* y *Aspergillus niger*. *Rev. Mex. Fitopatol.* 31:105-112.
- Cantrell C.L., K.K. Schrader, L.K. Mamonov, G.T. Sitpaeva, T.S. Kustova, C. Dunbar and D.E. Wedge. 2005. Isolation and Identification of Antifungal and Antialgal Alkaloids from *Haplophyllum sieversii*. *J. Agric. Food Chem.* 53:7741-7748.
- Cárdenas-Ortega, N.C., M.A. Zavala-Sánchez, J.R.C. Pérez-González and S. Pérez. 2005. Chemical composition and antifungal activity of essential oil of *Chrysactinia mexicana* Gray. *J. Agric. Food Chem.* 53:4347-4349.
- Carvalho, D.D.C., E. Alves, R. Barbosa-Camargos, D. Ferreira-Oliveira, J.R. Soares-Scolforoc, D. Antônio de Carvalho and T.R. Sâmia-Batista. 2011. Plant extracts to control *Alternaria alternata* in Murcott tangor fruits. *Rev. Iberoam. Micol.* 28:173-178.
- Castillo, F., D. Hernández, G. Gallegos, M. Méndez, R. Rodríguez, A. Reyes and C.N. Aguilar. 2010. In vitro antifungal activity of plant extracts obtained with alternative organic solvents against *Rhizotocnia solani* Kühn. *Ind. Crop Prod.*, 32:324-328.
- Castillo-Reyes, F., F.D. Hernández-Castillo, G. Gallegos-Morales, A. Flores-Olivas, R. Rodríguez-Herrera and

- C.N. Aguilar. 2015. Efectividad *in vitro* de *Bacillus* y polifenoles de plantas nativas de México sobre *Rhizoctonia-solani*. Rev. Mex. Ciencias Agríc. 6:549-562.
- Chávez-Quintal, P., T. González-Flores, I. Rodríguez-Buenfil and S. Gallegos-Tintoré. 2011. Antifungal Activity in Ethanolic Extracts of *Carica papaya* L. cv. Maradol Leaves and Seeds. Indian J. Microbiol. 51:54-60.
- Ciampi-Guillardi, M., C. Baldauf, A.P. Souza, G.J. Silva and L. Amorim. 2014. Recent introduction and recombination in *Colletotrichum acutatum* populations associated with citrus postbloom fruit drop epidemics in Sao Paulo, Brazil. Phytopathology. 104:769-778.
- Consolo, V.F., C.I. Mónaco, C.A. Cordo and G.L. Salerno. 2012. Characterization of novel *Trichoderma* spp. isolates as a search for effective biocontrollers of fungal diseases of economically important crops in Argentina. World J. Microbiol. Biotechnol. 28:1389-1398.
- Cotoras, M., C. Folch and L. Mendoza. 2004. Characterization of the Antifungal Activity on *Botrytis cinerea* of the Natural Diterpenoids Kaurenoic Acid and 3 β -Hydroxy-kaurenoic Acid. J. Agric. Food Chem. 52:2821-2826.
- Croplife. 2012. Value of fungicides in US crop production. Available online at: <http://www.croplifefoundation.org/upload/o>. Retrieved Feb. 17, 2014.
- Crous, P.W., G. Verkley, J. Groenewald and R. Samson. 2009. Fungal biodiversity. In: Fungal biodiversity. American Phytopathological Society (APS Press)).
- D'mello, J.P.F. and A.M.C. Macdonald. 2003. Fungal toxins as disease elicitors. p 255-294. In: J. Rose (ed.). Environmental Toxicology: current developments. Amsterdam: CRC Press.
- Damián-Badillo, L.M., R. Salgado-Garciglia, R.E. Martínez-Muñoz and M.M. Martínez-Pacheco. 2008. Antifungal Properties of Some Mexican Medicinal Plants. Open Nat. Prod. J., 1:27-33.
- Daoubi, M., R. Durán-Patrón, R. Hernández-Galán, A. Benharref, J.R. Hanson and I.G. Collado. 2006. The role of botrydiediol in the biodegradation of the sesquiterpenoid phytotoxin botrydial by *Botrytis cinerea*. Tetrahedron, 62:8256-8261.
- De Freitas, C.D.T., F.C. Sousa Nogueira, I.M. Vasconcelos, J.T. Abreu Oliveira, G. Barbosa Domont and M. Viana Ramos. 2011. Osmotin purified from the latex of *Calotropis procera*: Biochemical characterization, biological activity and role in plant defense. Plant Physiol. Bioch. 49:738-743.
- De León, M.A., A. Sáenz, D. Jasso-Cantu, R. Rodríguez, A. Pandey and C.N. Aguilar. 2013. Fermented *Flourensia cernua* extracts and their *in vitro* assay against *Penicillium expansum* and *Fusarium oxysporum*. Food Technol. Biotechnol. 51:233-239.
- De Lima-Guimarães, L.G., M. das Graças-Cardoso, P. Estevão de Sousa, J. de Andrade and S. Silveira-Vieira. 2011. Antioxidant and fungitoxic activities of the lemongrass essential oil and citral. Rev. Ciênc. Agron. 42:464-472.
- De Lucca, A.J. 2007. Hongos patógenos communes en la Agricultura y la Medicina. Rev. Iberoam. Micol. 24:3-13.
- De Menezes, H.D., A.C. Pereira, G.T.P. Brancini, H.C. de Leão, N.S. Massola Júnior, L. Bachmann, M. Wainwright, J.K. Bastos and G.U.L. Braga. 2014. Furocoumarins and coumarins photoinactivate *Colletotrichum acutatum* and *Aspergillus nidulans* fungi under solar radiation. J. Photoch. Photobio. B., 131:74-83.
- Dean, R., J.A. Van Kan, Z.A. Pretorius, K.E. Hammond-Kosack, A. Di Pietro, P.D. Spanu, J.J. Rudd, M. Dickman, R. Kahmann, J. Ellis and G.D. Foster. 2012. The Top 10 fungal pathogens in molecular plant pathology. Mol. Plant Pathol., 13(4):414-430.
- Del Castillo-Munera, J., M. Cardenas, A. Pinzon, A. Castaneda, A.J. Bernal and S. Restrepo. 2013. Developing a taxonomic identification system of *Phytophthora* species based on microsatellites. Rev. Iberoam. Micol. 30:88-95.
- Demirci, B., K.H.C. Baser, N. Tabanca and D.E. Wedge. 2006. Characterization of volatile constituents of *Haplopappus greenei* and studies on the antifungal activity against phytopathogens. J. Agric. Food Chem. 54:3146-3150.
- Díaz-Dellavalle, P., A. Cabrera, D. Alem, P. Larrañaga, F. Ferreira and M. Dalla-Rizza. 2011. Antifungal activity of medicinal plant extracts against phytopathogenic fungus *Alternaria* spp. Chil. J. Agr. Res., 71:231-239.
- Dos Santos Oliveira, M., and E. Badiale-Furlong. 2008. Screening of antifungal and antimycotoxigenic activity of plant phenolic extracts. World Mycotoxin J., 1:139-146.
- Duarte, Y., O. Pino and B. Martínez. 2014. Efecto de cuatro aceites esenciales sobre hongos asociados al manchado del arroz. Rev. Protección Veg. 29:62-65.
- Edin, E. 2012. Species specific primers for identification of *Alternaria solani*, in combination with analysis of the F129L substitution associates with loss of sensitivity toward strobilurins. Crop Prot., 38:72-73.
- Elamathi, R., R. Kavitha, P. Kamalakannan, T. Deepa and S. Sridhar. 2012. Preliminary phytochemical and antimicrobial studies on the leaf of *Ecbolium viride*. World. J. Pharm. Biol. Sci., 2(1):5-10.
- Espejo-Quispe, F., R. Espinoza-Ruiz, M. Puente-Isidró, R.C. Cupull-Santana, and M. Rodríguez-García. 2010. Efecto alelopático de *Tagetes erecta* y *Terminalia catappa* L sobre *Rhizoctonia solani* (Kühn). Centro Agrícola. 37:89-92.
- Eyre, C.A., M. Kozanitas and M. Garbelotto. 2013. Population dynamics of aerial and terrestrial populations of *Phytophthora ramorum* in a California forest under different climatic conditions. Phytopathology. 103:1141-1152.
- FAO. 2009. Regulation (EC) No 1107/2009 of the European Parliament and of the Council of 21 October 2009 concerning the placing of plant protection products on

- the market and repealing Council Directives 79/117/EEC and 91/414/EEC FAO council. (2009). Report of the twenty-first session of the committee on agriculture (COAG), Rome, April 22-25, 2009
- FAO. 2012. Mycotoxins. Available online at: <http://www.fao.org/food/food-safety-quality/a-z-index/mycotoxins/en/>. Retrieved August 20, 2012.
- FAO, IFAD, and WFP. 2013. The state of food insecurity in the world 2013. The multiple dimensions of food security. Rome, FAO
- Ferreira-Medeiros, J.G., A. Costa-Araujo-Neto, A. Silva de Medeiros, L. Cordeiro do Nascimento and E. Ursulino-Alves. 2013. Extratos vegetais no controle de patógenos em sementes de *Pterogyne nitens* Tul. Floresta e Ambiente. 20:384-390.
- Ferrucho, R.L., P.C. Ceresini, U.M. Ramirez-Escobar, B.A. McDonald, M.A. Cubeta and C. Garcia-Dominguez. 2013. The population genetic structure of *Rhizoctonia solani* AG-3PT from potato in the Colombian Andes. Phytopathology. 103:862-869.
- Fessehazion, B. and P. O. Olutiola. 1987. A proteolytic enzyme associated with the infection of cocoa beans by *Botryodiplodia theobromae*. FEMS Microbiol. Lett. 43:67-70.
- Finn, B.C., S. Vadala, A. Meraldi, J.E. Bruetman, J.V. Martinez, and P. Young. 2013. [Ergotism and HIV]. Medicina (B. Aires). 73:346-348.
- Fisher, A.J., T.R. Gordon and J.M. Ditomaso. 2005. Geographic distribution and diversity in *Claviceps purpurea* from salt marsh habitats and characterization of Pacific coast populations. Mycol. Res., 109:439-446.
- Flores Córdova, M.A., M.T. Martínez Damián, D. Nieto Ángel, J.E. Rodríguez Pérez, M.T. Colinas León and J. Martínez Solís. 2013. Reducción en la Germinación *in vitro* de conidios de *Alternaria alternata* aislada de *Eruca sativa* con Jugo de Brócoli. Rev. Mex. Fitopatol. 31:180-190.
- Freire, F.C., Z. Kozakiewicz and R.R. Paterson. 2000. Mycoflora and mycotoxins in Brazilian black pepper, white pepper and Brazil nuts. Mycopathologia. 149:13-19.
- Galvano, F., A. Piva, A. Ritieni and G. Galvano. 2001. Dietary strategies to counteract the effect of mycotoxins: a review. J. Food Prot. 64:120 – 131.
- Gambale, W., A. Purchio and J. Croce. 1976. Systematic isolation of the genus *Alternaria* in the atmosphere of the "Grande Sao Paulo" area (Brazil). Allergol. Immunopathol. (Madr), 4:139-144.
- Gamboa-Angulo, M.M., F. Alejos-Gonzalez, F. Escalante-Erosa, K. Garcia-Sosa, G. Delgado-Lamas and L.M. Peña-Rodriguez. 2000. Novel dimeric metabolites from *Alternaria tagetica*. J. Nat. Prod. 63:1117-1120.
- Gamboa-Angulo, M.M., F. Escalante-Erosa, K. Garcia-Sosa, F. Alejos-Gonzalez, G. Delgado-Lamas and L.M. Peña-Rodriguez. 2002. Natural zinnol derivatives from *Alternaria tagetica*. Isolation, synthesis, and structure-activity correlation. J. Agric. Food Chem. 50:1053-1058.
- Gamboa-Angulo, M.M., J. Cristóbal-Alejo, I.L. Medina-Baizabal, F. Chi-Romero, R. Méndez-González, P. Simá-Polanco and F. May-Pat. 2008. Antifungal properties of selected plants from the Yucatan peninsula, Mexico. World J. Microbiol. Biotechnol. 24:1955-1959.
- Garcia, R.A., F.C. Juliatti, K.A.G. Barbosa and T.A. Cassemiro. 2012. Antifungal activity of vegetable oils and extracts against *Sclerotinia Sclerotiorum*. Biosci. J. Uberlândia. 28:48-57.
- Garrido, C., M. Carbú, F.J. Fernández-Acero, V.E. González-Rodríguez and J.M. Cantoral. 2011. New Insights in the Study of Strawberry Fungal Pathogens. p. 24-39. In: Husaini Am and M. Ja. (ed.) Genes, Genomes and Genomics. Japan: Global Science Books.
- Garrido, C., F.J. Fernández-Acero, M. Carbú, V.E. González-Rodríguez, E. Liñeiro and J.M. Cantoral. 2012. Molecular Microbiology Applied to the Study of Phytopathogenic Fungi, Gel Electrophoresis - Advanced Techniques. (ed.) D.S. Magdeldin. InTech.
- Gomes, L.H., K.M.R. Duarte, F.G. Andrino, G.A. Leal Jr, L.M. Garcia, A. Figueira and S.P. de Lira. 2014. Alpha-Tocopherol against Witches' Broom Disease. Am. J. Plant Sci. 5:596-604.
- Gonzalez-Vera, A.D., J. Bernardes-De-Assis, M. Zala, B.A. McDonald, F. Correa-Victoria, E.J. Graterol-Matute and P.C. Ceresini. 2010. Divergence between sympatric rice- and maize-infecting populations of *Rhizoctonia solani* AG-1 IA from Latin America. Phytopathology. 100:172-182.
- Greco, M., A. Patriarca, L. Terminiello, V. Fernandez Pinto and G. Pose. 2012. Toxigenic *Alternaria* species from Argentinean blueberries. Int. J. Food Microbiol. 154:187-191.
- Gullino, M.L., P. Leroux and C.M. Smith. 2000. Uses and Challenges of novel compounds for plant disease control. Crop Prot. 19:1-11.
- Gunasekaran, M., and A.S. Ahuja. 1975. Effect of herbicides on mycelial growth of *Phymatotrichum omnivorum*. T. Brit. Mycol. Soc. 64:324-327.
- Gupta, S., and A.K. Dikshit. 2010. Biopesticides: An eco-friendly approach for pest control. J. Biopest. 3:186-188.
- Hardham, A.R. 2001. Cell biology of fungal infection of plants. p. 91-123. In: R. J. Howard and N. A. R. Gow (ed.) Biology of the Fungal Cell. Springer-Verlag Berlin Heidelberg.
- Harris, C.A., M.J. Renfrew and M.W. Woolridge. 2001. Assessing the risk of pesticide residues to consumers: recent and future development. Food Addit. Contam. 18:1124-1129.
- Hatcher, P.E., N.D. Paul, P.G. Ayres and J.B. Whittaker. 1994. The effect of a foliar disease (rust) on the development of *Gastrophysa viridula* (Coleoptera: Chrysomelidae). Ecol. Entomol. 19:349-360.
- Hernández, M., M. Murace, J. Ringuelet, I. Petri, D. Gallo and A. Arambarri. 2013. Effect of aqueous and alcohol extracts of *Phytolacca tetramera* (Phytolaccaceae) leaves on *Colletotrichum gloeosporioides* (Ascomycota). Bol. Soc. Argent. Bot. 48:201-209.

- Hernández, T., M. Canales, A.M. García, A. Duran, S. Meráz, P. Dávila and J.G. Ávila. 2008. Antifungal activity of the essential oils of two verbenaceae: *Lantana achyranthifolia* and *Lippia graveolens* of Zapotitlán de las Salinas, Puebla (México). Bol. Latinoam. Caribe Plant Med. Aromat. 7:203-207.
- Hernández-Silva, L., C. Piñón-Escobedo, H. Cano-Camacho, M.G. Zavala-Páramo, I. Acosta-Rodríguez and E. López-Romero. 2007. Comparison of fungal growth and production of extracellular pectin lyase activity by pathogenic and non-pathogenic races of *Colletotrichum lindemuthianum* cultivated under different conditions. Physiol. Mol. Plant P. 70:88-95.
- Hertel, S.C., H.D. Knöfel, R. Kramell and O. Miersch. 1997. Partial purification and characterization of a jasmonic acid conjugate cleaving amidohydrolase from the fungus *Botryodiplodia theobromae*. FEBS Letters. 407:105-110.
- Hope, J.H. and B.E. Hope. 2012. A Review of the Diagnosis and Treatment of Ochratoxin A Inhalational Exposure Associated with Human Illness and Kidney Disease including Focal Segmental Glomerulosclerosis. J. Environ. Public Health. 2012:835059
- International Energy Agency. 2014. Medium-term oil market report 2014.
- Inuma, T., S.A. Khodaparast and S. Takamatsu. 2007. Multilocus phylogenetic analyses within *Blumeria graminis*, a powdery mildew fungus of cereals. Mol. Phylogenet. Evol. 44:741-751.
- Iqbal, U. and T. Mukhtar. 2014. Morphological and Pathogenic Variability among *Macrophomina phaseolina* Isolates Associated with Mungbean (*Vigna radiata* L.) Wilczek from Pakistan. Sci. World J. I.D.950175:9
- Jaramillo, C.B.E., R.E. Duarte and W.D. Delgado. 2012. Bioactivity of essential oil from Colombian *Chenopodium ambrosioides*. Rev. Cubana Plant. Med. 17:54-64.
- Jasso de Rodríguez, D., D. Hernández-Castillo, J.L. Angulo-Sánchez, R. Rodríguez-García, J.A. Villarreal-Quintanilla and R.H. Lira-Saldivar. 2007. Antifungal activity *in vitro* of *Flourensia* spp. extracts on *Alternaria* sp., *Rhizoctonia solani*, and *Fusarium oxysporum*. Ind. Crop Prod. 25:111-116.
- Jasso de Rodríguez, D., R. Rodríguez-García, D. Hernández-Castillo, C.N. Aguilar-González, A. Saénz-Galindo, J.A. Villarreal-Quintanilla and L.E. Moreno-Zoccolotto. 2011. *In vitro* antifungal activity of extracts of Mexican Chihuahuan desert plants against postharvest fruit fungi. Ind. Crop Prod. 34:960-966.
- Johnson, D.A., G. Pimentel and F.M. Dugan. 2008. Cladosporium herbarum causes a Leaf Spot on Marshmarigold in Western North America. *Online Plant Health Progress* [Online]. Available: <http://www.plantmanagementnetwork.org/pub/php/research/2008/marsh/>.
- Kader, A.A. 2005. Increasing food availability by reducing postharvest losses of fresh produce. Acta Hort. 682:2169-2175.
- Kagale, S., T. Marimuthu, B. Thayumanavan, R. Nandakumar and R. Samiyappan. 2004. Antimicrobial activity and induction of systemic resistance in rice by leaf extract of *Datura metel* against *Rhizoctonia solani* and *Xanthomonas oryzae* pv. *oryzae*. Phys. Mol. Plant Pathol. 65:91-100.
- Knaak, N., L.D. da Silva, T.F. Andreis and L.M. Fiuza. 2013. Chemical characterization and anti-fungal activity of plant extracts and essential oils on the *Bipolaris oryzae* and *Gerlachia oryzae* phytopathogens. Australas. Plant Pathol. 42:469-475.
- Kobaisy, M., M.R. Tellez, C.L. Webber, F.E. Dayan, K.K. Schrader and D.E. Wedge. 2001. Phytotoxic and fungitoxic activities of the essential oil of Kenaf (*Hibiscus cannabinus* L.) leaves and its composition. J. Agric. Food Chem. 49:3768-3771.
- Kumar, V., M.S. Basu and T.P. Rajendran. 2008. Mycotoxin research and mycoflora in some commercially important agricultural commodities. Crop Prot. 27:891-905.
- Labbé, C., F. Faini, C. Villagrán, J. Coll and D.S. Rycroft. 2005. Antifungal and insect antifeedant 2-Phenylethanol Esters from the Liverwort *Balantiopsis cancellata* from Chile. Agric. Food Chem. 53:247-249.
- Lago, J.H., C.S. Ramos, D.C. Casanova, A. Morandim Ade, D.C. Bergamo, A.J. Cavalheiro, S. Bolzani Vda, M. Furlan, E.F. Guimaraes, M.C. Young and M.J.Kato. 2004. Benzoic acid derivatives from Piper species and their fungitoxic activity against *Cladosporium cladosporioides* and *C. sphaerospermum*. J. Nat. Prod. 67:1783-1788.
- Landero-Valenzuela, N., D. Nieto-Ángel, D. Téliz-Ortiz, R. Alatorre-Rosas, M. Orozco-Santos and C.F. Ortiz-García. 2013. Potencial antifúngico de extractos de cuatro especies vegetales sobre el crecimiento de *Colletotrichum gloeosporioides* en papaya (*Carica papaya*) en poscosecha. Rev. Venez. Cienc. Tecnol. Aliment. 4:047-062.
- Lara-Viveros, F.M., D. Nieto-Ángel, C. Nava-Díaz, G. Gutiérrez-Alonso, Ó.J. Ayala-Garay, L.A. Aguilar-Pérez, and T. Martínez-Damián. 2014. Efecto del glucorafano aislado de floretes de brócoli sobre la germinación de esporas de *Colletotrichum gloeosporioides*. Rev. Fitotec. Mex. 37:141-147.
- Latorre, B.A., I. Spadaro and M.E. Rioja. 2002. Occurrence of resistant strains of Botrytis cinerea to anilino pyrimidine fungicides in table grapes in Chile. Crop Prot. 21:957-961.
- Li, H.X. and C.L. Xiao. 2008. Baseline sensitivities to fludioxonil and pyrimethanil in *Penicillium expansum* populations from apple in Washington State. Postharvest Biol. Tec. 47:239-245.
- Li, S., Z. Zhang, A. Cain, B. Wang, M. Long and J. Taylor. 2005. Antifungal activity of camptothecin, trifolin, and hyperoside isolated from *Camptotheca acuminata*. J. Agric. Food Chem. 53:32-37.
- Lima de Sousa, L., S.C. Alburquerque de Andrade, A.J.A. Aguiar Athayde, C.E. Vasconcelos de Oliveira, C. Veríssimo de Sales, M. Suely Madruga and E. Leite de

- Souza. 2013. Efficacy of *Origanum vulgare* L. and *Rosmarinus officinalis* L. essential oils in combination to control posharvest pathogenic *Aspergilli* and autochthonous mycoflora in *Vitis labrusca* (table grapes). *Int. J. Food Microbiol.* 165:312-318.
- Lira-De León, K.I., M.V. Ramírez-Mares, V. Sánchez-López, M. Ramírez-Lepe, R. Salas-Coronado, N. F. Santos-Sánchez, R. Valadez-Blanco and B. Hernández-Carlos. 2014. Effect of crude plant extracts from some Oaxacan flora on two deleterious fungal phytopathogens and extract compatibility with a biofertilizer strain. *Front. Microbiol.* 5:1-10.
- Lourenco, V., A. Moya, F. Gonzalez-Candelas, I. Carbone, L.A. Maffia and E.S. Mizubuti. 2009. Molecular diversity and evolutionary processes of *Alternaria solani* in Brazil inferred using genealogical and coalescent approaches. *Phytopathology.* 99:765-774.
- Mabrouk, M.I. 2012. Synergistic and antibacterial activity of six medicinal plants used in folklore medicine in Egypt against *E. coli* O157:H7. *J. Appl. Sci. Res.* 8:1321-1327.
- Malbrán, I., C.A. Mourellos, J.R. Girotti, P.A. Balatti and G.A. Lori. 2014. Toxigenic Capacity and Trichothecene Production by *Fusarium graminearum* Isolates from Argentina and Their Relationship with Aggressiveness and Fungal Expansion in the Wheat Spike. *Phytopathology.* 104:357-364.
- Manandhar, H.K., H.J. Lyngs Jorgensen, S.B. Mathur and V. Smedegaard-Petersen. 1998. Suppression of Rice Blast by Preinoculation with Avirulent *Pyricularia oryzae* and the Nonrice Pathogen *Bipolaris sorokiniana*. *Phytopathology.* 88:735-739.
- Martín, M.D., S.Y. Ho, N. Wales, J.B. Ristaino and M.T. Gilbert. 2014. Persistence of the mitochondrial lineage responsible for the Irish potato famine in extant New World *Phytophthora infestans*. *Mol. Biol. Evol.*, 10.1093/molbev/msu086.
- Martins, J.A.S., E. Sagata, V.A. Santos and F.C. Juliatti. 2010. Evaluation of the effect of *Melaleuca alternifolia* oil on mycelial growth of phytopathogenic fungi *in vitro*. *Biosci. J., Uberlândia.* 27:49-51.
- Meepagala, K.M., K.K. Schrader, C.L. Burandt, D.E. Wedge and S.O. Duke. 2010. New class of algicidal compounds and fungicidal activities derived from a chromene amide of *Amyris texana*. *J. Agric. Food Chem.* 58:9476-9482.
- Meepagala, K.M., G. Sturtz and D.E. Wedge. 2002. Antifungal constituents of the essential oil fraction of *Artemisia dracunculus* L. Var. *dracunculus*. *J. Agric. Food Chem.* 50:6989-6992.
- Melo, V.M.M., I.K. Vasconcelos, V.M. Gomes, M. Da Cunha, A.A. Soares and J.T.A. Oliveira. 2005. A cotyledonary agglutinin from *Luetzelburgia auriculata* inhibits the fungal growth of *Colletotrichum lindemuthianum*, *Fusarium solani* and *Aspergillus niger* and impairs glucose-stimulated acidification of the incubation medium by *Saccharomyces cerevisiae* cells. *Plant Sci.* 169:629-639.
- Mendgen, K. and M. Hahn. 2002. Plant infection and the establishment of fungal biotrophy. *Trends Plant Sci.* 7:352-356.
- Monsálvez, M., N. Zapata, M. Vargas, M. Berti, M. Bittner and V. Hernández. 2010. Antifungal effects of *n*-hexane extract and essential oil of *Drimys winteri* bark against Take-All disease. *Ind. Crop Prod.* 31:239-244.
- Montes-Belmont, R. and A.M. Prados-Ligero. 2006. Influence of plant extracts on *Sclerotium cepivorum* development. *Plant Pathol. J.* 5:373-377.
- Moo-Koh, F.A., J. Cristóbal Alejo, A. Reyes-Ramírez, J.M. Tun-Suárez, R. Sandoval-Luna and J.A. Ramírez-Pool. 2014. Actividad *in vitro* del extracto acuoso del *Bonellia flammea* contra hongos fitopatógenos. *Agrociencia.* 48:833-845.
- Moulé, Y., F. Renauld, N. Darracq and C. Douce. 1982. DNA-protein cross-linking by the mycotoxin, botryodiplodin, in mammalian cells. *Carcinogenesis.* 3:211-214.
- Muñoz, G., P. Hinrichsen, Y. Brygoo and T. Giraud. 2002. Genetic characterisation of *Botrytis cinerea* populations in Chile. *Mycol. Res.* 106:594-601.
- Niño, J., O.M. Mosquera and Y.M. Correa. 2012. Antibacterial and antifungal activities of crude plant extracts from Colombian biodiversity. *Rev. Biol. Trop.* 60:1535-1542.
- Nogueira, J.H.C., E. Gonçalves, S.R. Galleti, R. Facanali, M.O.M. Marques and J.D. Felício. 2010. *Ageratum conyzoides* essential oil aflatoxin suppressor of *Aspergillus flavus*. *Int. J. Food Microbiol.* 137:55-60.
- Noreña-Ramírez, D.A., O.J. Velásquez-Ballesteros, E. Murillo-Perea, J.J. Méndez-Arteaga, J.F. Solanilla-Duque and W. Murillo-Arango. 2014. Assessment of *Phaseolus vulgaris* L and *Vigna unguiculata* (L.) Walp leaves for antifungal metabolites against two bean fungal pathogens *Colletotrichum lindemuthianum* and *Phaeoisariopsis griseola*. *Afr. J. Biotechnol.* 13:1657-1665.
- O'Donnell, K., H.C. Kistler, B.K. Tacke and H.H. Casper. 2000. Gene genealogies reveal global phylogeographic structure and reproductive isolation among lineages of *Fusarium graminearum*, the fungus causing wheat scab. *Proc. Natl. Acad. Sci. U.S.A.* 97:7905-7910.
- Oliva, A., K.M. Meepagala, D.E. Wedge, D. Harries, A.L. Hale, G. Aliotta and S.O. Duke. 2003. Natural fungicides from *Ruta graveolens* L. leaves, including a new quinolone alkaloid. *J. Agric. Food Chem.* 51:890-896.
- Ortiz-Núñez, Y., I. Spengler-Salabarría, I.G. Collado and R. Hernández-Galán. 2007. Sesquiterpenes from the wood of *Juniperus lucayana*. *Phytochemistry.* 68:2409-2414.
- Osorio, E., M. Flores, D. Hernández, J. Ventura, R. Rodríguez and C.N. Aguilar. 2010. Biological efficiency of polyphenolic extracts from pecan nuts shell (*Carya illinoensis*), pomegranate husk (*Punica granatum*) and creosote bush leaves (*Larrea tridentata* Cov.) against plant pathogenic fungi. *Ind. Crop Prod.* 31:153-157.
- Otálvaro, F., J. Nanclares, L.E. Vásquez, W. Quiñones, F. Echeverri, R. Arango and B. Schneider. 2007.

- Phenalenone-Type compounds from *Musa acuminata* var. "Yangambi km 5" (AAA) and their activity against *Mycosphaerella fijiensis*. J. Nat. Prod. 70:887-890.
- Pane, C., R. Spaccini, A. Piccolo, F. Scala and G. Bonanomi. 2011. Compost amendments enhance peat suppressiveness to *Pythium ultimum*, *Rhizoctonia solani* and *Sclerotinia minor*. Biol. Control. 56:115-124.
- Pavón-Moreno, M.A., I. Gonzalez-Alonso, R. Martin-De Santos and T. Garcia-Lacarra 2012. [The importance of genus *Alternaria* in mycotoxins production and human diseases]. Nutr. Hosp. 27:1772-1781.
- Pedras, M.S.C., M.G. Sarwar, M. Suchy and A.M. Adio. 2006. The phytoalexins from cauliflower, caulilexins A, B and C: Isolation, structure determination, syntheses and antifungal activity. Phytochemistry. 67:1503-1509.
- Pekkarinen, A.I. and B.L. Jones. 2003. Purification and Identification of Barley (*Hordeum vulgare* L.) Proteins that inhibit the Alkaline Serine Proteinases of *Fusarium culmorum*. J. Agric. Food Chem. 51:1710-1717.
- Pelegri, P.B., L.R. Farias, A.C.M. Saude, F.T. Costa, C. Bloch Jr., L.P. Silva, A.S. Oliveira, C.E.M. Gomes, M.P. Sales and O.L. Franco. 2009. A Novel Antimicrobial Peptide from *Crotalaria pallida* seeds with activity against human and phytopathogens. Curr. Microbiol. 59:400-404.
- Peraza-Sánchez, S.R., E.O. Chan-Che and E. Ruiz-Sánchez. 2005. Screening of Yucatecan Plant Extracts to control *Colletotrichum gloeosporioides* and Isolation of a New Pimarene from *Acacia pennatula*. J. Agric. Food Chem. 53:2429-2432.
- Perelló, A., M. Gruhlke and A.J. Slusarenko. 2013 Effect of garlic extract on seed germination, seedling health, and vigour of pathogen-infested wheat. J. Plant Prot. Res. 53:317-323.
- Pérez-Láinez, D., R. García-Mateos, R. San Miguel-Chávez, M. Soto-Hernández, E. Rodríguez-Pérez, and G. Kite. 2008. Bactericidal and fungicidal activities of *Calia secundiflora* (Ort.) Yakovlev. Z. Naturforsch. 63c:653-657.
- Pérez-Martin, J. and C. De Sena-Tomas. 2011. Dikaryotic cell cycle in the phytopathogenic fungus *Ustilago maydis* is controlled by the DNA damage response cascade. Plant Signal Behav. 6:1574-1577.
- Perrone, G., G. Stea, F. Epifani, J. Varga, J.C. Frisvad and R.A. Samson. 2011. *Aspergillus niger* contains the cryptic phylogenetic species A. awamori. Fungal Biol. 115:1138-1150.
- Pfohl-Leszkowicz, A. and R.A. Manderville. 2007. Ochratoxin A: An overview on toxicity and carcinogenicity in animals and humans. Mol. Nutr. Food Res. 51:61-99.
- Phillipson, B. 2012. Quarantine Procedures for Working with Fungal Plant Pathogens. p. 119-137. In: Lane Ch., Beales P., Hughes K. (ed.) Fungal Plant Pathogens.ed. CABI, South Asia.
- Pimentel, D. 2006. Soil erosion: a food and environmental threat. Environ. Dev. Sustain. 8:119-137.
- Pimentel, D., C. Harvey, P. Resosudarmo, K. Sinclair, D. Kurz, M. McNair, S. Crist, L. Sphpritz, L. Fitton, R. Saffouri and R. Blair. 1995. Environmental and economic costs of soil erosion and conservation benefits. Science. 267:1117-1123.
- Pineda, R., S. Vizcaino, C.M. García, J.H. Gil and D.L. Durango. 2012. Chemical Composition and Antifungal Activity of *Piper auritum* Kunth and *Piper holtonii* C. DC. Against phytopathogenic fungi. Chil. J. Agr. Res. 72:507-515.
- Pollock, G.A., C.E. Disabatino, R.C. Heimsch and D.R. Hilblink. 1982. The subchronic toxicity and teratogenicity of alternariol monomethyl ether produced by *Alternaria solani*. Food Chem. Toxicol. 20:899-902.
- Poole, G.J., R.W. Smiley, C. Walker, D. Huggins, R. Rupp, J. Abatzoglou, K. Garland-Campbell and T.C. Paulitz. 2013. Effect of climate on the distribution of *Fusarium* spp. causing crown rot of wheat in the Pacific Northwest of the United States. Phytopathology. 103:1130-1140.
- Population Reference Bureau. 2014. 2014 World Population Data Sheet. Washington D.C.
- Prandini, A., S. Sigolo, L. Filippi, P. Battilani and G. Piva. 2009. Review of predictive models for *Fusarium* head blight and related mycotoxin contamination in wheat. Food Chem. Toxicol. 47:927-931.
- Prieto, J.A., O.J. Patiño, W.A. Delgado, J.P. Moreno and L.E. Cuca. 2011. Chemical Composition, Insecticidal, And Antifungal Activities of Fruit Essential Oils of Three Colombian *Zanthoxylum* Species. Chil. J. Agr. Res. 71:73-82.
- Pupo-Blanco, Y.G., D. Kalombo-Bicayi, L. Herrera-Isla, D.I. Malheiros de Mendonca and B. Vargas-Batis. 2011. Effect of plant extracts on growth and spore germination of *Alternaria solani* (E. & M.) J. & G. under *in vitro* conditions. Rev. Iberoam. Micol. 28(1):60.
- Qin, X.F. and F.B. Kong. 2006. Prospect and situation of biopesticides. J. Anhui Agric. Sci. 16:4024-4057.
- Queiroz, S.C.N., C.L. Cantrell, S.O. Duke, D.E. Wedge, V.K. Nandula, R.M. Moraes and A.L. Cerdeira. 2012. Bioassay-directed isolation and identification of phytotoxic and fungitoxic acetylenes from *Conyza Canadensis*. J. Agric. Food Chem. 60:5893-5898.
- Quiroga, E.N., A.R. Sampietro and M.A. Vattuone. 2001. Screening antifungal activities medicinal plants. J. Ethnopharmacol. 74:89-96.
- Ramezani, M., W.T. Shier, H.K. Abbas, J.L. Tonos, R.E. Baird and G.L. Sciumbato. 2007. Soybean charcoal rot disease fungus *Macrophomina phaseolina* in Mississippi produces the phytotoxin (-) -botryodiplodin but no detectable phaseolinone. J. Nat. Prod. 70:128-129.
- Ramírez Aguayo, H.E., G. Virgen-Calleros, J.D.J. Vargas-Radillo, E. Salcedo-Pérez and L. Barrientos-Ramírez. 2015. Actividad antimicrobiana in vitro de extractos de hoja de *Guazuma ulmifolia* Lam. contra fitopatógenos. Rev. Mex. Cienc. Forest. 6:114-124.
- Ramírez-Chávez, E., L. Lucas-Valdez, G. Virgen-Calleros and J. Molina-Torres. 2000. Fungicidal activity of

- afinnin and *Heliopsis longipes* crude root extract on two *Sclerotium* species. *Agrocienci.* 34:207-215.
- Richard, J., Payne, G., Eds, Desjardins, A., Maragos, C., Norred, W., and Pestka, J. 2003. "Mycotoxins: risks in plant, animal and human systems," in CAST Task Force Report. Council for Agricultural Science and Technology Ames, IA. 101-103.
- Rodriguez-Maturino A., R. Troncoso-Rojas, A. Sánchez-Estrada, D. González-Mendoza, E. Ruiz-Sanchez, R. Zamora-Bustillos, C. Ceceña-Duran, O. Grimaldo-Juarez and M. Aviles-Marin. 2015. Antifungal effect of phenolic and carotenoids extracts from chiltepin (*Capsicum annum* var. *glabriusculum*) on *Alternaria alternata* and *Fusarium oxysporum*. *Rev. Argent. Microbiol.* 47:72-77.
- Rodríguez-Rajo, F.J., I. Iglesias and V. Jato. 2005. Variation assessment of airborne *Alternaria* and *Cladosporium* spores at different bioclimatical conditions. *Mycol. Res.* 109:497-507.
- Rostás, M., M. Simon and M. Hilker. 2003. Ecological cross-effects of induced plant responses towards herbivores and phytopathogenic fungi. *Basic Appl. Ecol.* 4:43-62.
- Rouissi, W., L. Ugolini, C. Martini, L. Lazzeri and M. Mari. 2013. Control of postharvest fungal pathogens by antifungal compounds from *Penicillium expansum*. *J. Food Prot.* 76:1879-1886.
- Sá, R.A., A.C.C. Argolo, T.H. Napoleão, F.S. Gomes, N.D.L. Santos, C.M.L. Melo, A.C. Albuquerque, H.S. Xavier, L.C.B.B. Coelho, L.W. Bieber and P.M.G. Paiva. 2009. Antioxidant, *Fusarium* growth inhibition and *Nasutitermes corniger* repellent activities of secondary metabolites from *Myracrodruon urundeuva* heartwood. *Int. Biodeter. Biodegr.* 63:470-477.
- Salah, M.A., E. Bedir, N.J. Toyang, I.A. Khan, M.D. Harries and D.E. Wedge. 2003. Antifungal clerodane diterpenes from *Macaranga monandra* (L.) Muell. Arg. (Euphorbiaceae). *J. Agric. Food Chem.* 51:7607-7610.
- Saleh, A.A., H.U. Ahmed, T.C. Todd, S.E. Travers, K.A. Zeller, J.F. Leslie and K.A. Garrett. 2010. Relatedness of *Macrophomina phaseolina* isolates from tallgrass prairie, maize, soybean and sorghum. *Mol. Ecol.* 19:79-91.
- Sampietro, D.A., C.G. Diaz, V. Gonzalez, M.A. Vattuone, L.D. Ploper, C.A. Catalan and T.J. Ward. 2011. Species diversity and toxigenic potential of *Fusarium graminearum* complex isolates from maize fields in northwest Argentina. *Int. J. Food Microbiol.* 145:359-364.
- San Gupta, R., R.R. Chandran and P.V. Divekar. 1966. Botryodiplodin--a new antibiotic from *Botryodiplodia theobromae*. II. Production, isolation and biological properties. *Indian J. Exp. Biol.* 4:152-153.
- Sánchez-García, C., M. Cruz, E.L. Martín, M. Leyva-Mora, M. Cruz-Martín, Y. Alvarado-Capo, M. Acosta-Suárez, B. Roque and M. Pérez. 2008. Actividad antifúngica del aceite esencial de *Cymbopogon nardus* para el control de *Macrophomina phaseolina*. *Centro Agrícola.* 35:83-86.
- Sánchez-Martín, J., D. Rubiales and E. Prats. 2011. Resistance to powdery mildew (*Blumeria graminis* f.sp. *avenae*) in oat seedlings and adult plants. *Plant Pathol.* 60:846-856.
- Santana, G.M.S., L.P. Albuquerque, D.A. Simões, L.C.B.B. Coelho, P.M.G. Paiva and N.B. Gusmão. 2009. Isolation of lectin from *Opuntia ficus-indica* cladodes. *Acta Hort.* 811:281-286.
- Schinke, C. and J.C. Germani. 2012. Screening Brazilian *Macrophomina phaseolina* isolates for alkaline lipases and other extracellular hydrolases. *Int. Microbiol.* 15:1-7.
- Schoebel, C.N., J. Stewart, N.J. Gruenwald, D. Rigling and S. Prospero. 2014. Population History and Pathways of Spread of the Plant Pathogen *Phytophthora plurivora*. *PLoS One*, 9, e85368.
- Sharma, K. and N.M. Meshram. 2006. Bioactivity of essential oils from *Acorus calamus* Linn. and *Syzygium aromaticum* Linn. against *Sitophilus oryzae* Linn. In stored wheat. *Biopest. Int.* 2:144-152.
- Siddiqui, F.A. and T. Gulzar. 2003. Tetracyclic triterpenoids from the leaves of *Azadirachta indica* and their insecticidal activities. *Chem. Pharm. Bull.* 51:415-417.
- Somma, S., G. Pose, A. Pardo, G. Mule, V.F. Pinto, A. Moretti and A.F. Logrieco. 2011. AFLP variability, toxin production, and pathogenicity of *Alternaria* species from Argentinean tomato fruits and puree. *Int. J. Food Microbiol.* 145:414-419.
- Svetaz, L., A. Tapia, S.N. López, R.L.E. Furlán, E. Petenatti, R. Pioli, G. Schmeda-Hirschmann and S.A. Zacchino. 2004. Antifungal chalcones and new caffeic acid esters from *Zuccagnia punctata* acting against soybean infecting fungi. *J. Agric. Food Chem.* 52:3297-3300.
- Tabanca, N., B. Demirci, S.L. Crockett, K.H.C. Başer and D.E. Wedge. 2007. Chemical Composition and antifungal activity of *Arnica longifolia*, *Aster hesperius*, and *Chrysanthamnus nauseosus* essential oils. *J. Agric. Food Chem.* 55:8430-8435.
- Tagami, O.K., M.D.G. Gasparin, K.R.F. Schwan-Estrada, M.E. da Silva Cruz, A.T. Itako, J.B. Tolentino Júnior, L.M. de Moraes and Stangarlin, J.R. 2009. Fungitoxicity of *Bidens pilosa*, *Thymus vulgaris*, *Lippia alba* and *Rosmarinus officinalis* in the *in vitro* development of phytopathogens fungi. *Semin-Cienc. Agrar.* 30:285-294.
- Tanaka, K., Y. Sago, Y. Zheng, H. Nakagawa and M. Kushiro. 2007. Mycotoxins in rice. *Int. J. Food Microbiol.* 119:59-66.
- Taylor, J.L. and M.S.C. Pedras. 1995. Horizontal transfer in the phytopathogenic fungal genus *Leptosphaeria* and host-range expansion. *Trends Microbiol.* 3:202-206.
- Taylor, R.J., B. Salas, G.A. Secor, V. Rivera and N.C. Gudmestad. 2002. Sensitivity of North American Isolates of *Phytophthora erythrosptica* and *Pythium ultimum* to Mefenoxam (Metalaxyl). *Plant Dis.* 86:797-802.
- Teixeira de Freitas C.D., L.L. de Souza Lopes, L.M. Beltrami, R.S. Basílio de Oliveira, J.T. Abreu Oliveira and M.V. Ramos. 2011. Osmotin from *Calotropis procera* latex: New insights into structure and antifungal properties. *Biochim. Biophys. Acta.* 10:2501-2507.
- Thangavel, T., R. Steven Tegg and C.R. Wilson. 2014. Resistance to multiple tuber diseases expressed in

- somaclonal variants of the potato cultivar russet burbank. Sci. World J. ID:417697.
- Tittlemier, S.A., D. Gaba and J.M. Chan. 2013a. Monitoring of *Fusarium trichothecenes* in Canadian cereal grain shipments from 2010 to 2012. J. Agric. Food Chem. 61:7412-7418.
- Tittlemier, S.A., M. Roscoe, R. Trelka, D. Gaba, J.M. Chan, S.K. Patrick, M. Sulyok, R. Krska, T. Mckendry and T. Grafenhan. 2013b. *Fusarium* damage in small cereal grains from Western Canada. 2. Occurrence of fusarium toxins and their source organisms in durum wheat harvested in 2010. J. Agric. Food Chem. 61:5438-5448.
- Umpierrez-Failache, M., G. Garmendia, S. Pereyra, A. Rodriguez-Haralambides, T.J. Ward and S. Vero. 2013. Regional differences in species composition and toxigenic potential among *Fusarium* head blight isolates from Uruguay indicate a risk of nivalenol contamination in new wheat production areas. Int. J. Food Microbiol. 166:135-140.
- Uppalapati, S.R., C.A. Young, S.M. Marek and K.S. Mysore. 2010. Phymatotrichum (cotton) root rot caused by *Phymatotrichopsis omnivora*: retrospects and prospects. Mol. Plant Pathol. 11:325-334.
- Urdaneta, L., M.E. Sanabria, D. Rodríguez, G. Ettiene and M. Pérez. 2015. Flavonoides en hojas de *Gliricidia sepium* y su efecto *in vitro* sobre *Colletotrichum acutatum*. Bol. Centro Invest. Biol. 48:115-130.
- Vaartaja, O. 1974. Inhibition of *Pythium ultimum* in molecular fractions from gel filtration of soil extracts. Can. J. Microbiol. 20:1273-1280.
- Valenzuela-Cota, D.F., G.V. Buitimea-Cantúa, E.C. Rosas-Burgos, F.J. Cinco-Moroyoqui, M.S. Yépiz-Gómez, M.O. Cortez-Rocha, M. Plascencia-Jatomea and A. Burgos-Hernández. 2014. Efecto antifúngico de los extractos de la planta *Jacquinia macrocarpa* en el desarrollo de *Aspergillus flavus*, *A. parasiticus* y *Fusarium verticillioides*. Rev. Mex. Micol. 39:1-11.
- Vargas-Arispuru, I., A. Contreras-Valenzuela and M.A. Martínez-Téllez. 2009. Lignans from *Larrea tridentate* (creosote Bush) as fungal β -1,3-glucanase inhibitors. Pestic. Biochem. Phys. 94:60-63.
- Veloz-García, R., R. Marín-Martínez, R. Veloz-Rodríguez, R. Rodríguez-Guerra, I. Torres-Pacheco, M.M. González-Chavira, J.L. Anaya-López, L. Guevara-Olvera, A.A. Feregrino-Pérez, P. Loarca-Piña and R.G. Guevara-González. 2010. Antimicrobial activities of cascálope (*Caesalpinia cacalaco*) phenolics-containing extract against fungus *Colletotrichum lindemuthianum*. Ind. Crop Prod. 31:134-138.
- Vituro, C.I., J.R. de la Fuente and M.S. Maier. 2004. 5-Methylcoumaranones from *Mutisia friesiana* and their bioactivity. J. Nat. Prod. 67:778-782.
- Vogt, V., D. Cifuentes, C. Tonn, L. Sabini and S. Rosas. 2013. Antifungal activity *in vitro* and *in vivo* of extracts and lignans isolated from *Larrea divaricata* Cav. against phytopathogenic fungus. Ind. Crop Prod. 42:583-586.
- Wogan, G.N. 1999. Aflatoxin as a human carcinogen. Hepatology. 30:573-575.
- Wu, H., D. Wang, J. Shi, S. Xue and M. Gao. 2010. Effect of the complex of zinc (II) and cerium (IV) with chitosan on the preservation quality and degradation of organophosphorus pesticides in Chinese jujube (*Zizyphus jujuba* Mill. cv. Dongzao). J. Agric. Food Chem. 58:5757-5762.
- Wyand, R.A. and J.K. Brown. 2003. Genetic and forma specialis diversity in *Blumeria graminis* of cereals and its implications for host-pathogen co-evolution. Mol. Plant Pathol. 4:187-198.
- Xu, Y.Z. 2008. The application of biological pesticides and industrial development measures. Chin. Agr. Sci. Bull. 24:402-404.
- Yanes, L.R., P.I. Torres, R.G. Guevara-Gonzales, M.I. Hernandez-Zul, J.A. Quijano-Carranza and E. Rico-Garcia. 2012. The effect of climate change on plant diseases. Afr. J. Biotechnol. 11:2417-2428.
- Yang, H.C., J.S. Haudenschild and G.L. Hartman. 2014. *Colletotrichum incanum* sp. nov., a curved-conidial species causing soybean anthracnose in USA. Mycologia. 106:32-42.
- Yoganathan, K., L.K. Yang, C. Rossant, Y. Huang, S. Ng, M.S. Butler and A.D. Buss. 2004. Cochlioquinones and epi-cochlioquinones: antagonists of the human chemokine receptor CCR5 from *Bipolaris brizae* and *Stachybotrys chartarum*. J. Antibiot. 57:59-63.
- Zamora-Natera, F., P. García-López, M. Ruiz-López, J. Ruiz Moreno, M. Pedrosa and M. Muzquiz. 2008. Composition of Alkaloids in seeds of *Lupinus mexicanus* (Fabaceae) and antifungal evaluation of the alkaloid extract. In: J.A. Palta and J.B. Berger (eds). Lupins for Health and Wealth, eds. J.A. Palta and J.B. Berger Proceedings of the 12th International Lupin Conference, 14-18 Sept. 2008, Fremantle, Western Australia. International Lupin Association, Canterbury, New Zealand. ISBN 0-86476-153-8.
- Zhang, Y.B. 2009. Analyze the importance of pesticides based on world's needs on grain and agricultural development. World Pestic. 31:1-3.